



THE INTERNATIONAL ASSOCIATION OF  
**GEOMAGNETISM AND AERONOMY**  
INTERNATIONAL UNION OF GEODESY AND GEOPHYSICS



IAGA-ITALIA

## Resoconto di attività 2011-2014

Prof. U. Villante

Delegato Italiano IAGA

# International Association of Geomagnetism and Aeronomy (IAGA)

---

## IAGA Activities in Italy (2011-2014)

([www.iagaitalia.it](http://www.iagaitalia.it))

IAGA activities in Italy are currently developed by several universities as well as by mayor Scientific Institutions such as Istituto Nazionale di Geofisica e Vulcanologia (INGV), Istituto Nazionale di Astrofisica (INAF) and Consiglio Nazionale delle Ricerche (CNR).

The present document, organized on the basis of IAGA Divisions and Interdisciplinary Commissions, summarizes the principal achievements, the participation to international programs and the most relevant programs in which the Italian Scientific community is involved.

IAGA Community in Italy is organized according to the following scheme:

Italian Correspondent: Prof. U. Villante, University of L'Aquila

Vice Correspondent: Dr. L. Vigliotti, CNR – Istituto di Scienze Marine (ISMAR), Bologna

|  |   |
|--|---|
| Division I:<br>(pag. 3)                  | “Internal Magnetic Field” (Coordinator: Dr. L. Vigliotti – CNR - ISMAR, Bologna)            |
| Division II:<br>(pag. 8)                 | “Aeronomic Phenomena” (Coordinator: Dr. B. Zolesi – INGV - Roma)                            |
| Division III:<br>(pag. 13)               | “Magnetospheric Phenomena” (Coordinator: Dr. G. Consolini – INAF/IAPS, Roma)                |
| Division IV:<br>(pag. 17 )               | “Solar Wind and Interplanetary Magnetic Field” (Coordinator: Dr R. Bruno – INAF/IAPS, Roma) |
| Division V:<br>(pag. 21)                 | “Geomagnetic Observatories, Surveys and Analyses” (Coordinator: Dr A. Meloni – INGV, Roma)  |
| Interdivisional Commission:<br>(pag. 25) | “History” (Coordinator: Dr A. De Santis – INGV, Roma)                                       |
| Interdivisional Commission:<br>(pag.25 ) | “Education and Outreach” (Coordinator: Prof. F. Berrilli – Università Tor Vergata, Roma)    |

Meetings of the Italian IAGA Community have been held on:

- October 2011 – IAGA/Italy General Assembly
- June 2013 – IAGA /Italy Meeting - Future Projects and Programs

As summarized in the following, the Italian IAGA Community organized several conferences and international schools.

The Italian IAGA Community also played a major role in the establishment of the National Group for Space Weather (Oct. 2013).

## DIVISION I

### “Internal Magnetic Field” (Coord.: L. Vigliotti)

#### A) Institutions involved in research activity

1. Istituto Nazionale di Geofisica e Vulcanologia, INGV-Roma.
2. Centro Interuniversitario di Magnetismo Naturale (CIMaN-ALP ALPINE LABORATORY OF PALEOMAGNETISM University of Milano, Torino, Parma, Roma3, Chieti-Pescara and Urbino).
3. Department of Physics of the University of Bari.
4. Department of Physics of the University of Camerino.
5. University of Naples.
6. Istituto di Scienze Marine ISMAR-CNR, Bologna.
7. Istituto di Astrofisica Spaziale e Planetologia/INAF Roma

#### B) Scientific Report

##### B1) Main Scientific Themes

Interdisciplinary studies concerning the registration of the Earth magnetic field, magnetic properties of rocks/sediments and studies of the electromagnetic signal focused on different subjects including:

**Magnetostratigraphy.** Using Earth's magnetic field reversal as a method of dating rocks.

**Earth's magnetic field.** Structure and paleosecular variations of Earth's magnetic field from marine/lacustrine sediments and archaeological data (Archaeomagnetic dating).

**Environmental Magnetism.** Paleoclimatic and paleoceanographic reconstructions based on magnetic properties of sediments. Study of atmospheric particulate even trapped in the Antarctic Ice.

**Tectonics.** Tectonic evolution of the Mediterranean region as well as areas in other continents: Asia, Africa, South America.

**Volcanology.** Use of paleomagnetism and rock magnetism to understand volcanic processes

**Sedimentology.** Depositional patterns of turbidites rocks.

- **Paleomagnetism, Magnetic polarity and Secular Variation record of the magnetic field**

Paleomagnetic studies achieved important results in the reconstruction of the dynamics and timing of polarity reversals of the Earth's magnetic field. The study of Sagnotti et al. (2014) of lake sediments from the Basin of Sulmona showed that the transition of the magnetic polarity during the reversal would take place in a time comparable to that of human life. This study could have major implications not only for understanding the behavior of the Earth's magnetic field. With regard to the integrated stratigraphy and magnetostratigraphy, researches focused on several sections ranging in time from Cretaceous to Pleistocene. Some of them are candidates for the definition of different Global Stratotype Section and Point (GSSP). Early Miocene (Burdigalian) sections were studied in the Conero, Iblei as well as in oceanic cores. Additional GSSP focused on the Paleogene (Lutetian, Thanetian, Bartonian and Selandian) and in particular on stratigraphic series sampled at Gorrondatxe and Zumaia in the Basque countries (Spain). Additional studies on land took into account sections from the Anatolian Plateau (Turkey), New Zealand, the Piedmont Tertiary Basin and Pleistocene sequences from the Padania Plain. Measurements on marine sediments include the upper part of the stratigraphic succession recovered in the IODP expedition 317 (Canterbury Basin).

Several investigations have been carried out on the paleosecular variations (PSV) of the Earth magnetic field recorded by sediments (Adriatic Sea, Ionian Sea, Barents Sea) and

volcanic rocks (Azores, Capo Verde, Pantelleria), with a re-evaluation of the Italian historical geomagnetic catalogue. The analysis of different cores from the Barents Sea, off the coast of Svalbard, has allowed the reconstruction of the Holocene PSV at high northern latitudes, providing important experimental constraints for models of the geomagnetic field, and allowing the development of an age model for detailed reconstruction of climatic variations. Magnetostratigraphic investigations include also classic sites inhabited by hominid in Puglia (Pirro Nord) and at Ceprano.

- **Environmental Magnetism**

Magnetic parameters were used to investigate the paleoclimatic and paleoceanographic history recorded by marine and lacustrine sediments in different environments. Results have been obtained on sediments representing extreme climatic conditions such as greenhouse episodes during the Eocene-Paleocene and the Messinian salinity crisis, but also from young sediments. In the framework of the International Continental Drilling Program (ICDP) “Paleovan”, 150 meters of Late Quaternary sediments collected at Site 5034 in the Lake Van (Turkey) were analyzed for a paleomagnetic/rock magnetic study carried out in collaboration with the University of Florida at Gainesville (USA).

Magnetic data from Antarctic and periantarctic drillings have been summarized to reconstruct the climatic evolution of this region during the Cenozoic.

Application to studies of environmental pollution, especially by particulate matter were carried out with the aim to determine the contribution of the ultrafine magnetic fraction (diameter less than 30 nm) in atmospheric dust, characterized by superparamagnetic properties.

Electrical and electromagnetic methods were used in various geodynamic environments at different scales of investigation in Italy, Morocco and especially in volcanic areas. The targets focused on the continuous monitoring of the magnetotelluric field, development of techniques for the characterization of the magnetotelluric signal, study of the seismic-electromagnetic signals.

- **Geodynamic Reconstructions**

With regard to applications to geodynamics, paleomagnetic and magnetic anisotropy were applied to reconstruct the tectonic evolution of the Italian peninsula, several Mediterranean regions and the Andes. Studies in the Croton basin provided new original data for the geodynamic mechanisms responsible of the Plio-Pleistocene deformation of the Calabrian block. In the Andes region, new studies have been conducted in the Eastern Cordillera of Colombia to constrain the mechanisms for the genesis of this mountain chain. New paleomagnetic data were obtained from Paleozoic rocks in Sardinia as well as from the Neogene of Central Iran. The anisotropy of magnetic susceptibility was used to reconstruct the Pleistocenian deformation field in the upper valley of the Tiber.

- **Magnetic Properties of Rocks and Archeomagnetism**

An increasing number of studies focused on the applications of magnetic parameters to volcanic rocks. The emplacement temperatures and depositional mechanism of piroclastic and volcanic units were analyzed to characterize different volcanic eruptions (Breccia di Commenda, Brown Tuff, Ferrar Dolerites, Pollena, Pompei, Colima, El Chichon).

Archeomagnetic studies took into account furnaces found at Kato Achaia (Grecia), Fontanetto Po, Spilamberto, Portugal, and on ceramics from Skala Sotiros, Archontiko, Tempi, Paros, and Santorini (Greece). Studies for determination the firing temperature of ceramic artefacts were carried out in collaboration with the University of Bari and the Archaeological Office of different regions (Piemonte, Emilia-Romagna) and also the Canton of Bern in Switzerland. The origin and composition of Neolithic obsidian as well as Roman and Medieval bricks were investigated in collaboration with the National Institute of Metrological Research to

reconstruct the provenance and the commercial roads in the Mediterranean region during the ancient times.

Specific research of the remanent magnetization carried out by magnetic minerals of biogenic origin in marine sediments gave new insights on their influence in the acquisition and retention of the magnetic properties. Additional results focused on peculiar magnetic properties of different species of magnetotactic bacteria.

## **B2) Main Research Projects/Programmes**

- Magnetic models and analyses in the Project “Magnetic Homing. PRIN
- Earth Magnetic Field measurements, Modelling and Seismicity – EM3S Italy-Albania. Bilateral project DS3F “Deep Sea and Subseafloor Frontier”
- Satellite, ground and seafloor measurement for earthquake pattern recognition”. SAGA-4-EPR
- Paleomagnetic and Magnetic properties of the Lake Van sediments (PALEOVAN PROJECT; ICDP SITE 5034): Paleoclimatic and environmental reconstruction of the last 500000 years. Short-Term mobility” program.
- Acque di fusione glaciale, plumiti e morene recessionali allo sbocco della Fossa di Storfjorden durante la deglaciazione della Calotta Glaciale del Mare di Barents (MELTSTORM)
- Studio multidisciplinare dei sedimenti glaciomarini depositi nel Mare di Ross (Antartide) negli ultimi 50 Ka: informazioni sulle fluttuazioni dell'estensione dei ghiacci nel corso della transizione glaciale-interglaciale
- Rapporti Temporali tra Attività Vulcanica e Sedimentazione nell'Area McMurdo Sound - Ross Ice Shelf: Implicazioni Tettoniche e Paleoclimatiche. ANDRILL- ANTArctic DRILLing .
- Paleomagnetic Analyses of Drifting and Tectonic Rotations of Central Iran. Progetto nell'ambito del Consorzio internazionale DARIUS (2009-2011).
- Magnetostratigraphy and stratigraphy of Miocene to Present lacustrine basins in central Anatolia. Collaborative Research Project Vertical Anatolian Movement Project (VAMP), Progetto TOPOEUROPE (ESF) (2008-2012).
- Osservazioni di terreno e sperimentali per la stima della pericolosità vulcanica Valutazione della pericolosità vulcanica in termini probabilistici. Progetto Vulcanologico DPC-INGV. Task 2.
- Genesi e differenziazione dei magmi in relazione all'ambiente geodinamico ed alle caratteristiche petrologiche e geochemiche delle loro sorgenti: implicazioni per l'evoluzione del sistema convergente Africa-Europa. PRIN.
- Indagini ad alta risoluzione per la stima della pericolosità e del rischio sismico nelle aree colpite dal terremoto del 6 aprile 2009. FIRB.
- EPOS (European Plate Observing system) preparatory phase.
- CORIBAR-IT: Dinamica glaciale e deposizione associata all'ultimo ritiro glaciale del Mare di Barents nordoccidentale: un progetto a 5-nazioni per perforare l'Artico con il sistema MeBo. PNRA
- Reconstructing 2500 years of environmental change at the periphery of Rome: Integrating paleoecology and socioeconomic history to understand human response to climate.
- ROSSLOPE II: Dinamica sedimentaria passata ed attuale nel Mare di Ross: un approccio multidisciplinare allo studio della scarpata continentale. PNRA
- Il "GSSP" (Global Stratigraphic Section and Point) del Piano Burdigaliano: il tassello mancante all'intervallo Neogenico della Scala del Tempo Geologico
- The potential of AMS (Anisotropy of Magnetic Susceptibility) studies in polyphase deformed rocks.
- REGENA project “Holocene and Pleistocene recovery of geomagnetic secular variation in the North Atlantic (Azores and Cape-Verde archipelagos and Portugal mainland): Geomagnetic and volcanological implications”. FCT Project (Portugal)
- MED-SUV (MEDiterranean SUPersite Volcanoes), FP7

- Project ENI Val d'Agri: subcontract with the Institute for the Methodologies for Environmental Analysis of the CNR (CNR\_IMAA) of Potenza.
- Geochimica, geodinamica, e cinematica delle placche nel Mar Rosso. PRIN
- Integrated assessment of magnetotelluric and magnetospheric monitoring in seismic areas of L'Aquila and the Val d'Agri for the identification, quantification and interpretation of possible modifications of the electrical resistivity in the earth's crust

### **B3) Funding Agencies**

- MIUR PRIN Magnetic models and analyses in the Project “Magnetic Homing”
- MIUR PRIN Valutazione integrata del monitoraggio magnetotellurico e magnetosferico nelle aree sismiche de L'Aquila e della Val d'Agri ai fini della identificazione, quantificazione e interpretazione di eventuali modificazioni della resistività elettrica nella crosta terrestre
- MIUR PRIN Geochimica, geodinamica, e cinematica delle placche nel Mar Rosso.
- Italian Foreign Office Bilateral project Italy-Albania 2012-2013.
- European Project DS3F “Deep Sea and Subseafloor Frontier” 2010-2012
- Italian Foreign Office SAGA-4-EPR Project of Excellence Italy-China 2010-2012
- CNR “Short-Term mobility program” "Paleomagnetic and Magnetic properties of the Lake Van sediments (PALEOVAN PROJECT; ICDP SITE 5034): Paleoclimatic and environmental reconstruction of the last 500000 years"
- PNRA Acque di fusione glaciale, plumiti e morene recessionali allo sbocco della Fossa di Storfjorden durante la deglaciazione della Calotta Glaciale del Mare di Barents (MELTSTORM)
- PNRA Studio multidisciplinare dei sedimenti glaciomarini depositi nel Mare di Ross (Antartide) negli ultimi 50 Ka: informazioni sulle fluttuazioni dell'estensione dei ghiacci nel corso della transizione glaciale-interglaciale
- PNRA ANDRILL-ANTarctic DRILLing - Rapporti Temporali tra Attività Vulcanica e Sedimentazione nell'Area McMurdo Sound - Ross Ice Shelf : Implicazioni Tettoniche e Paleoclimatiche
- Provincia Autonoma di Bolzano - Alto Adige
- CNR. Progetto Bandiera RITMARE.
- NSF-USA
- ENI

### **B4) Organization of Conferences and Workshop of international relevance**

Italian researchers from different institutions were convenors of important international conferences:

- M. Mattei (Roma-3) co-convenor of the J-V07 symposia “Using Paleomagnetism to Understand Volcanic Processes” IUGG, 2011, Melbourne, Australia.
- P. Macrì (INGV) Co-convenor of the EMRP1 Open session on “Geomagnetism and Paleomagnetism” EGU General Assembly 2011, Vienna, Austria.
- L. Sagnotti (INGV) Convenor della sessione EMRP3.1 “Open session on Rock magnetism and Paleomagnetism”, EGU General Assembly 2014; Vienna, Austria.
- A. De Santis (INGV) Convenor session EMRP2.1 “Open Session on Geomagnetism“, EGU General Assembly 2014, Vienna, Austria.
- E. Tema (Università di Torino). Convenor della sessione EMRP3, “Archaeological material, volcanic rocks and sediments as past geomagnetic field recorders: New data, methods and applications.” EGU General Assembly 2014; Vienna, Austria.
- F. Florindo (INGV) Convenor sessione “Environmental Magnetism in Climate, Stratigraphy and Environmental Studies”. AGU, San Francisco, Dicembre, 2014.

- A. De Santis (INGV) co-organiser of the International Scientific Spring (ISS-2014), Islamabad (Pakistan) March 2014.

### **B5) Prizes**

- G. Currenti (University of Catania) IAGA Medal at the IUGG meeting (Melbourne 2011)
- A. De Santis (INGV) has been appointed as Vice-President of the Earth Magnetism and Rock Physics (EMRP) Division of the European Geoscience Union (EGU) in the years 2012-2014.

## **C) Goals, priorities and plans for future activities**

### **C1) Main Scientific Themes**

- Development of an integrated network of research infrastructures, in particular rock physics labs, in the framework of the EPOS project
- Integrated magnetostratigraphy, as a high-resolution correlation and dating tool for marine and continental stratigraphic sequences even in sequences representing GSSP
- Paleomagnetism applied to geodynamics; new experimental data and insights for understanding the evolution of the central Mediterranean and the genesis of complex orogens
- Environmental magnetism as an original proxy for the reconstruction of paleoenvironmental and paleoclimatic changes, with particular emphasis on the Mediterranean and both polar regions
- Environmental magnetism, as an original tool to investigate particulate matter air pollution
- Reconstruction of paleosecular variation of the geomagnetic field for the Holocene and relative paleointensity for the Pleistocene. Application to correlation and dating of geologic events with implication for the evaluation of volcanic and seismic risks.

### **C2) Main Research Projects/Programmes**

- European Plate Observing System (EPOS)  
Funding agency: European Community
- Studio multidisciplinare dei sedimenti glaciomarini depositi nel Mare di Ross (Antartide) negli ultimi 50 Ka: informazioni sulle fluttuazioni dell'estensione dei ghiacci nel corso della transizione glaciale-interglaciale  
Funding agency: PNRA
- NEXTDATA un sistema nazionale per la raccolta, conservazione, accessibilità e diffusione dei dati ambientali e climatici in aree montane e marine  
Funding agency: MIUR
- Indagini ad alta risoluzione per la stima della pericolosità e del rischio sismico nelle aree colpite dal terremoto del 6 aprile 2009  
Funding agency: MIUR
- Reconstructing 2500 years of environmental change at the periphery of Rome: Integrating paleoecology and socioeconomic history to understand human response to climate  
Funding agency: NSF
- ARctic: present Climatic change and pAst extreme events (ARCA) MIUR
- PRIN Geochimica, geodinamica, e cinematica delle placche nel Mar Rosso.  
Funding Agency: MIUR.

## **DIVISION II**

### **“Aeronomic Phenomena” (Coord.: B. Zolesi)**

#### **A) Institutions involved in research activity**

1. Istituto Nazionale di Geofisica e Vulcanologia, Roma ;
2. Istituto dei Sistemi Complessi, Consiglio Nazionale delle Ricerche, Firenze
3. “Abdus Salam “ International Centre for Theoretical Physics, Trieste.

#### **B) Scientific Report**

##### **B1) Main Scientific Themes**

##### **Istituto Nazionale di Geofisica e Vulcanologia**

- **HF and L-band ionospheric monitoring**

Presently, ionospheric vertical soundings are performed in two ionospheric observatories in Italy, Rome (41.8° N, 12.5° E) and Gibilmanna (37.9° N, 14.0° E), and one in the Italian Antarctic base M. Zucchelli. Considerable cooperation has been activated with the Argentinean colleagues to install and operate the Italian ionospheric station AIS in Tucuman (26.9° S, 294.6° E), Argentina.

Hourly systematic measurements of the critical frequency of the F2 layer, foF2, recorded at the Rome and Gibilmanna ionospheric observatories, along with the hourly quiet time reference values of foF2, foF2<sub>QT</sub>, were considered around the periods of minimum and maximum solar activity over the years 1976-2008, to study the ionospheric variability over Rome and Gibilmanna.

Power Virtual Height measurements (PVH) of radio echoes reflected from the ionosphere acquired at the Rome ionospheric observatory have been used to study the multipath fading variations through time of the ionospheric channel on a temporal scale from 0.5 to 256 s.

INGV is managing a network of high rate (50 Hz) sampling GNSS (Global Navigation Satellite Systems) receivers at polar, equatorial and mid-latitudes. The network now counts instruments in Antarctica (at Mario Zucchelli Station and at Concordia), in Arctic (at Svalbard Islands), in the Mediterranean area (Lampedusa and Rome, Italy; Crete, Greece) and in Argentina (Tucuman). Through a network funded by different projects INGV cooperates also to the operation of similar receivers in Brazil. Such observations are used to investigate the irregular behaviour of the ionosphere causing scintillation, a diffractive and refractive effect on the trans-ionospheric signals transmitted by GNSS satellites. INGV has achieved a robust expertise in modelling the scintillation and has activated several international collaborations in the field.

- **Ionospheric morphology and modelling**

Unusual nighttime impulsive electron density enhancements that are rarely observed at low latitudes on a wide region of South America, under quiet and medium/high geomagnetic conditions, were analyzed. A detailed analysis of isoheight ionosonde plots suggests that traveling ionospheric disturbances (TIDs) caused by gravity wave (GW) propagation could play a significant role in causing the phenomenon both for quiet and for medium/high geomagnetic activity; in the latter case however a recharging of the fountain effect, due to electric fields penetrating from the magnetosphere, joins the TID propagation and plays an as much significant role in causing impulsive electron density enhancements. An atypical nighttime spread-F structure appearing on ionograms at or above the F2 trace, near the crest of the ionospheric equatorial ionization anomaly (EIA) region, was investigated. The phenomenon appeared to be the signature of MSTIDs propagating above or at F2-layer peak that generates an extra ionospheric structure just above or at the F2-layer, during nighttime. A



comparative study of spread-F signatures over five low-latitude sites was performed. The study confirmed that the dynamics and the physical processes responsible for these phenomena are actually complicated. In fact, the features that arise from the investigation are different, depending both on the longitude sector and on the hemisphere. A spread-F and GPS scintillation statistics of occurrence under the southern crest of the equatorial ionospheric anomaly was carried out. The most striking features coming from the study were a systematic correspondence between SSF (Strong Range Spread-F) and scintillations, and a possible correlation between  $S_4$  and FSF (Frequency Spread-F) peaks at the terminator. Assuming that both  $fE_s$  and  $h'E_s$  trends are influenced by the atmospheric tides, the height–time–intensity (HTI) technique was applied to deeply investigate how these waves control the  $E_s$  dynamics. The HTI study, along with a fast Fourier transform analysis, show that a well-defined semidiurnal periodicity characterizes the  $E_s$  layer dynamics most accurately in June and July, while in August and September the daytime semidiurnal periodicity becomes weaker and the role of the diurnal periodicity is consequently highlighted. The behavior of the critical frequency of the F2 region of the ionosphere ( $f_oF_2$ ) and the height of the maximum density of free electrons in F2 region ( $h_mF_2$ ) over Tucumán (26.91S, 294.61E), during the deep solar minimum occurred in 2008–2009. Data used were compared with those obtained at solar minimum observed in 1975–1976. An ionospheric service was implemented in the Mediterranean area around Nicosia (Cyprus) in terms of: plots in real-time of the ionospheric characteristics  $f_oF_2$  and  $f_{min}$  recorded by the digisonde installed at Nicosia; nowcasting and long-term maps of  $f_oF_2$ ,  $M(3000)F_2$ , and MUF (with the transmitting point located at Nicosia) over a regional area around Nicosia.

A short-term ionospheric forecasting empirical regional model (IFERM) to predict the state of the critical frequency of the F2 layer ( $f_oF_2$ ) over Europe during moderate, disturbed, and strongly geomagnetic conditions, as well as a short-term ionospheric forecasting empirical regional model to predict  $M3000F_2$  over Europe during not quiet geomagnetic conditions plus a method for  $f_oF_2$  short-term (1–24 h) forecast using both historical and real-time  $f_oF_2$  observations over European stations, EUROMAP model, has been developed. The predictions of  $M3000F_2$  are obtained correcting the monthly median field of  $M3000F_2$  with the Bradley - Dudeney formula that depends on the short-term predictions of  $f_oF_2$  provided by IFERM model. Hourly systematic measurements of the critical frequency of the F2 layer,  $f_oF_2$ , recorded at the Rome (Italy, 41° 8' N, 12° 5' E) and Gibilmanna (Italy, 37° 6' N, 14° 0' E) ionospheric observatories, along with the hourly quiet time reference values of  $f_oF_2$ ,  $f_oF_{2QT}$ , were considered around the periods of minimum and maximum solar activity over the years 1976–2008, to study the ionospheric variability over Rome and Gibilmanna. A comparative study between the sporadic E layers over the ionospheric stations of Rome and Gibilmanna has been carried out using hourly systematic measurements of the highest frequency reflected by the  $E_s$  layer ( $f_oE_s$ ) recorded during the period January 1976 - July 2008 at the Rome and Gibilmanna ionospheric observatories. Power Virtual Height measurements (PVH) of radio echoes reflected from the ionosphere acquired at the Rome ionospheric observatory have been used to study the multipath fading variations through time of the ionospheric channel on a temporal scale from 0.5 to 256 s. A three-dimensional (3-D) IRI-SIRMUP-P (ISP) model of electron density of the ionosphere was developed and tested over several periods characterized by very disturbed geomagnetic conditions. The three-dimensional (3-D) electron density representation of the ionosphere computed by the assimilative IRI-SIRMUP-P (ISP) model was used in connection with IONORT (IONospheric Ray-Tracing), a software application for calculating a 3-D ray-tracing in the ionospheric medium. For testing the reliability of IONORT-system, several comparisons between measured oblique ionograms over the radio link between Rome (41.8\_N, 12.5\_E) and Chania (35.7\_N, 24.0\_E), and synthesized oblique ionograms from the IONORT – system have been performed. In the framework of the project ESA Space Weather Network -IV3 project European ionosonde and neutron monitor services, two original software for the achievement of maps of  $f_oF_2$  and  $M3000F_2$  extending from 34°

N to 80° N in latitude and from -10° W to 40° W in longitude are under development. In the framework of the project CIFS (Cyprus Ionospheric Forecasting Service), long term and nowcasting maps of the maximum usable frequency (MUF) were generated over the east region of Mediterranean extending from 30° N to 40° N in latitude and from 20° W to 45° W in longitude. Studies concerning the effects of Small Scale Disturbance (SSD) on signal fading and coherence times of the ionosphere have been conducted on the basis of vertical ionospheric soundings at 3 MHz performed at the Rome ionospheric observatory. A short-term forecasting regional model (STFRM), based on the integrated geomagnetic index  $ap(\tau)$  has been developed to predict M(3000)F2 over the European area during moderate, disturbed, and very disturbed geomagnetic conditions. The possible relationship between thunderstorms and the ionospheric sporadic E Layer over Rome has been investigated using the superposed epochs analysis. Studies about the possible influence of solar cycle on the critical frequencies of the mid-latitude sporadic E layer have been carried out using the Continuous Wavelet Transform technique, Fast Fourier Transform, and linear regression analyses. In order to obtain a three dimensional electron density mapping of the ionosphere up to three hours in advance, studies concerning the achievement of short term forecasting of electron density profiles over some stations located in European area are in progress.

Seasonal (Winter/Summer) and solar cycle NmF2 variations as well as summer saturation effect in NmF2 have been analyzed using Millstone Hill ISR daytime observations. A new mechanism (qualitative) to explain the December anomaly in NmF2 is proposed. A new method has been developed to retrieve neutral temperature  $T_n$  and composition [O], [N<sub>2</sub>], [O<sub>2</sub>] from electron density profiles in the daytime mid-latitude F2-region under both quiet and disturbed conditions.

Other themes of main interest are: empirical and theoretical modeling of ionospheric scintillations to forecast and mitigate the ionospheric corruptions on GNSS signals; scientific understanding of the cause-effect mechanism leading from the ionospheric irregularities formation to the scintillation effects; structuring and accessibility of the ionospheric data for real-time and post processing use. The empirical and theoretical modeling of ionospheric scintillations is based on the experience matured on the analysis of the data owned by INGV and its partners. The modelling is addressed to forecast and to mitigate the ionospheric corruptions on GNSS signals. The experimental activity allowed to advance the scientific understanding of the cause-effect mechanism leading from the ionospheric irregularities formation to the scintillation effects. The significant amount of data collected since 2003 has been structured and made accessible to registered users via the eSWua Database: [www.eswua.ingv.it](http://www.eswua.ingv.it).

#### **Istituto dei Sistemi Complessi, Consiglio Nazionale delle Ricerche, Firenze**

During the years 2010-2014 CNR's activity in the field of near-Earth space research has been concentrated mainly in the study of the dynamics of complex/irregular plasmas and near-Earth space imaging/observation. CNR team conducts the project named ISIS, ESA-CNR about the observation of the Earth's plasmasphere.

#### **Abdus Salam “ International Centre for Theoretical Physics, Trieste.**

The Telecommunications/ICT for Development Laboratory (T/ICT4D) (formerly Aeronomy and Radiopropagation Laboratory) of the Abdus Salam International Centre for Theoretical Physics has concentrated its ionosphere research and applications work in the following lines: Earth and Mars Ionosphere modelling, data ingestion and assimilation in models, use of radio occultation for ionosphere characterization, lower latitudes ionosphere, ionosphere effects in GNSS operations at range and position domains. The work has been carried out in collaboration with groups of European and African universities.

### **B2/3) Main Research Projects/Programmes and funding Agencies**

- CIGALA project “Concept for Ionospheric Scintillation Mitigation for Professional GNSS in Latin America” FP7, Collaborative projects, GALILEO 2008GSA, 2010-2012.
- ESPAS: “Near-Earth Space Data Infrastructure for e-Science” a FP7 project.
- Royal society Project: in collaboration with University of Nottingham and INGV funded by the Royal Society (UK): "Original and Novel Solutions to Counter GNSS Ionospheric Scintillation Effects", 2010-2012;
- Research Project IDIPOS "Italian Data Base Infrastructure for Polar Observation Science", funded by National Program of Antarctic Research (PNRA).
- Research Project 2009/B03 "Upper Atmosphere Observations and Space Weather", 2009-2012, funded by National Program of Antarctic Research (PNRA).
- SWING : (Short Wave Critical infrastructure Network based on new Generation prediction Tools) EU Directorate Justice, Freedom and Security. 2012- 2013
- TELEDIFE :Training courses and Ionospheric radio propagation prediction; Ministry of Italian Defense
- TRANSMIT project (Training Research and Applications Network to Support the Mitigation of Ionospheric Threats), 2011-2015, FP7 Marie Curie-ITN.
- MIMOSA, GINESTRA, MEDSTEC 2012-2013, ESA Alcantara initiatives to survey the ionospheric monitoring capabilities over South America, Africa, Asia.
- Funding Agencies : FP7, ESA, PNRA, Ministry of Italian Defense.
- CALIBRA (Countering GNSS high Accuracy applications Limitations due to Ionospheric disturbances in BRAzil) FP7, 2012-2014.
- MIMOSA2, ERICA 2014-2016, Continuation of ESA Alcantara initiatives.
- MISW (Mitigation of space weather threats to GNSS services), FP7, 2014-2016.
- DemoGRAPE, PNRA, 2014-2016.
- CIFS ,Cyprus Ionospheric Forecasting Service, 2012-2014

### **B4) Organization of conferences and Workshop of international relevance**

- M.Pezzopane as convenor of the session “S4: Space Weather effects on HF and trans-ionospheric radio wave propagation” at the 10<sup>th</sup> European Space Weather Week, Antwerp, Belgium, 18-22 November 2013.
- M.Pezzopane as member of the “Scientific Organizing Committee” of the session “C1.1: Recent Advances in Equatorial, Low- and Mid-Latitude Mesosphere, Thermosphere and Ionosphere Studies” at the COSPAR (Committee on SPace Research) scientific assembly, Mysore, India, 14-22 July 2012.

### **B5) Prizes**

- M. Pezzopane: 2013. First prize for the best communication presented at the XCIX National Congress of the Italian Physical Society, held in Trieste from 23 to 27 September 2013, Section 4a “Geophysics, Environmental Physics, Oceanography”; title of the communication “Unusual nighttime impulsive enhancements of electron density characterizing the low-latitude ionosphere: phenomenology and possible mechanisms of triggering”.

### **C) Goals, priorities and plans for future activities**

- The study of TEC and ionospheric scintillation will be further advanced to improve the scientific understanding and to increase the performance of the forecasting and mitigation modeling. The results of new studies on the ionospheric impact on the SAR observations and on the regional assessment of the polar scintillations will be presented in the next 2 years.
- Different spectral analyses will be employed to investigate the tidal and planetary wave periodicities imprinted in the following two main characteristics of the sporadic (Es) layer: the top frequency (ftEs) and the lowest virtual height (h'Es). Studies are planned to verify the

influence of the solar activity on the Es layer. The behavior of the ionospheric plasma in correspondence of the anomalous and prolonged minimum of solar activity between cycles 23 and 24 will be investigated.

## **DIVISION III**

### **“Magnetospheric Phenomena” (Coord.: G. Consolini)**

#### **A) Institutions involved in research activity**

1. Institute of Space Astrophysics and Planetology of the Istituto Nazionale di Astrofisica (INAF-IAPS)
2. Istituto Nazionale di Geofisica e Vulcanologia
3. Department of Physics of the University of L'Aquila

#### **B) Scientific Report**

##### **B1) Main Scientific Themes**

##### **Physics of the Magnetosphere.**

The magnetospheric studies in Italy are mainly focused on 1) the interaction between the solar wind and the planetary magnetospheres (mainly the Earth's magnetosphere), 2) the Earth's magnetospheric dynamics, 3) the processes responsible for the plasma transport in the magnetospheric regions, and 4) the Earth's magnetosphere-ionosphere coupling.

During 2010-2011 a particular interest has been devoted to the study of acceleration plasma processes, such as magnetic reconnection, which are responsible for the enhancement of plasma transfer and transport in the inner magnetospheric regions, and to the study of ULF waves and pulsations, of the emergence of complexity and the role of turbulence and intermittence in the global magnetospheric dynamics. These studies are mainly based on magnetic field and plasma data recorded on-board of magnetospheric spacecrafts (e.g., CLUSTER, Geotail, Double-Star,...) and on ground-based geomagnetic and ionospheric observatories. Using data coming from Cluster and Double Star TC-1, it has been possible to study the interaction of the solar wind with the terrestrial magnetosphere. In particular, research has been performed concerning processes like reconnection, interaction of interplanetary shocks with the magnetosphere. Such research has been funded by ASI in the framework of the contract I/023/09/0 which ended in June 2012. Using detailed Double Star observations of proton distribution functions and magnetic field data around a Flux Transfer Event (FTE) at the terrestrial magnetopause, useful information on the FTE generation mechanism have been obtained. These observations have suggested that two X lines active at the magnetopause are probably the generation mechanism of the observed FTE confirming the Lee and Fu FTE model of multiple reconnection X lines rather than the patchy reconnection proposed in the Russell and Elphic model. In the last two years a detailed study of the magnetic and electric field fluctuations at the kinetic scale has been done. This study allowed to reveal the emergence of different scaling features at scales below the ion-inertial length during magnetic reconnection.

Special attention has been paid to the study of the magnetic storms and substorms, which are two important manifestations of the interaction between the solar wind and the Earth's magnetic field. The relationship between these two phenomena is a key point in the study of the Sun-Earth connection and despite numerous studies conducted over the past years, there are still problems that are not completely solved. In this context, during the period 2011-2012, the relationship between these two classes of magnetospheric phenomena has been studied through an approach based on the information theory. In detail, the dependence of the information transfer on the conditions of the solar wind has been investigated. It has been found that the flow of information between these two classes of magnetospheric phenomena depends on the global level of magnetic activity. The influence (in terms of information flows)

of substorms on storms is maximum in the case of moderate/intermediate geomagnetic activity level while the driving of storms by substorms seems to be less effective during intense magnetic storms. Furthermore, studying the multiscale features of some large geomagnetic storms by applying the empirical mode decomposition technique it has been shown that there is an increase of dynamical complexity and multi-scale properties for intermediate geomagnetic activity levels. This increase seems to reflect the influence of substorms to storms that is maximum during these periods. In contrast, during low and high geomagnetic activity levels, the geomagnetic field fluctuations seem to be a consequence of a very stochastic dynamics, similar to the global dynamics that characterize a Markovian nonequilibrium relaxation process.

Using geomagnetic field data collected on the ground, the magnetospheric dynamics has been investigated analyzing the low frequency geomagnetic pulsations and the daily variation of the Earth's magnetic field. In detail, the analysis of magnetic data collected by geomagnetic observatories placed at high latitude all located within the polar cap but at different distance from the auroral oval (Antarctic and Arctic) has allowed analyzing the diurnal variation of the magnetic field. The results have shown that the diurnal variation strongly depends on the position within the polar cap, both regarding its shape and its dependence on season, solar wind and magnetospheric parameters. At the same time, a statistical analysis of low frequency (1-5 mHz) pulsations, at different Antarctic stations situated along the same geomagnetic parallel (80S) within the polar cap, has allowed to study the azimuthal propagation of the observed signals; the results have shown that the pulsations are originated preferably in two different regions: around local geomagnetic noon and around local geomagnetic midnight; they are interpreted in term of cusp-related and geomagnetic tail-related phenomena, respectively; these signals are well coherent between station spaced up to 5 hours in magnetic local time. The analysis, conducted also using data from Cluster and Geotail, has suggested that waves can transmit across the magnetopause flanks and tail lobes and, propagating along the outermost field lines, reach the ground at polar latitudes. The ULF magnetometer stations, which have allowed to achieve positive results on the dynamic magnetospheric studies, have been installed in a new area. Indeed, as responsible of the Work Package "Retrieval of equatorial plasma mass densities by magnetometer arrays and cross-calibration" an Italian group of researchers has participated to the activities connected to the FP7 Project PLASMION ("A new, ground based data-assimilative model of the Earth's plasmasphere - a critical contribution to Radiation Belt modeling for Space Weather purposes", 2011-2014). In the framework of this project, a new latitudinal array of 25 ULF magnetometer stations (EMMA) extending from Italy to Finland (including SEGMA) has been created. A code has been developed to deduce in quasi-real-time the equatorial plasma mass density at different L-shells using the field line eigen-frequencies automatically determined from the EMMA array and the real-time solar wind parameters from the ACE satellite which are necessary for modelling the magnetospheric field by the Tsyganenko model. Data coming from ULF magnetometer stations have been used to analyze the possibility that electromagnetic signals can precede the occurrence of an earthquake. In particular, several papers have been dedicated to the possible occurrence of electromagnetic signals preceding the L'Aquila earthquake (6 April, 2009) mostly in the ULF range. The results do not show, in general, any clear evidence for ULF precursors.

In the year 2013 the INAF-IAPS group started an international collaboration in the framework of the activities connected to the FP7 Project STORM (Solar system plasma turbulence, observations, intermittency and multifractals) coordinated by Dr. Marius Echim of the BIRA-IASB (Brussels, Belgium). This project aims to investigate the relevance and the features of intermittent turbulence in several interplanetary and near-planetary environments, among which Earth's and planetary magnetosheaths from the magnetohydrodynamic scales to the kinetic ones. Furthermore, the project is also devoted to the analysis of the intermittent and

multifractal properties of short- and long-term magnetospheric dynamics as it results from the analysis of the features of geomagnetic indices time series. Preliminary results (communicated to international meetings - papers in preparation) have clearly shown how the critical nature of the Earth's magnetosphere depends on the solar cycle.

The Italian community (INGV and INAF-IAPS groups) started a study of the turbulent and scale-invariance features of the magnetic field variations on short temporal and small spatial scales as observed by the novel ESA mission Swarm, launched in the late 2013. The aim of these studies is the understanding of the influence of the different magnetospheric disturbances (magnetic storms and substorms) and currents on the ionospheric turbulence for a better understanding and modeling of the inner magnetospheric dynamics and magnetosphere-ionosphere coupling. The first results, relative to the investigation of the high latitude scaling features of the horizontal component of the magnetic field, allowed for the characterization of the turbulent spectra as a function of the geomagnetic activity level (AE-index) in the polar regions.

Finally, the Italian community is involved in the studies concerning the global solar wind - magnetosphere - ionosphere coupling and the inter-hemispheric magnetic conjugacy. Such studies are performed throughout the measurements provided by the SuperDARN network. In relation to this activity, in late 2012 – early 2013 the Dome C East radar has been successfully installed in the framework of a French-Italian co-operation (LPC2E-CNRS and INAF/IAPS). The context of solar wind - magnetosphere - ionosphere coupling includes also the analysis of SI (Sudden Impulses) events, which have been measured both at geostationary orbit and on the ground. This analysis, allowing careful comparisons with theoretical models of the magnetospheric field representation, has been useful to distinguish the contribution of magnetospheric and ionospheric current systems to the magnetospheric and geomagnetic perturbations.

## **B2) Main Research Projects/Programmes**

- **PNRA-PEA (U. Villante, resp.)**  
Title: ULF pulsations, magnetospheric dynamics and Space Weather aspects at polar latitudes (2011-2013).
- **FP7-SPACE-2010.2.3-1 PLASMON (M. Villante, resp.)**  
Title: A new, ground based data-assimilative modeling of the Earth's plasmasphere - a critical contribution to Radiation Belt modeling for Space Weather purposes (2011-2014).
- **FP7-SPACE-2012.2.1-01 STORM (M. Echim - BIRA, Brussels, Belgium-, resp., G. Consolini Italian group resp.)**  
Title: Solar system plasma turbulence: observations, intermittency and multifractals (2013-2015).
- **PNRA-PEA (L. Cafarella, resp.)**  
Title: Osservazioni di geomagnetismo ed elettromagnetismo in Antartide.
- **PNRA-PEA (M.F. Marcucci, resp.)**  
Title: Studio bipolare di fenomeni magnetosferici con SuperDARN ed osservazioni ottiche e magnetiche (2014-2016).

## **B3) Funding Agencies**

- Italian National Programme for Antarctica Research (PNRA)
- Italian Space Agency (ASI)
- European Commission (FP7-Space Call)

## **B5) Prizes**

In the year 2011 Prof. Umberto Villante received the "Antonio Feltrinelli" Prize on Astronomy, Geodesy, Geophysics and applications, assigned by the Italian Accademia dei Lincei, for his relevant scientific activity in the field of interplanetary space and magnetospheric physics.

## **C) Goals, priorities and plans for future activities (next two years, max 1 page)**

In these next two years the scientific activities of the previous period will continue. In particular, the Italian groups will devote to:

- The studies of the interaction of solar wind structures with the Earth's magnetosphere, with a special emphasis to the influence of the magnetospheric and ionospheric current systems in the manifestation of magnetic storms and substorms, as well as sudden impulses of the magnetospheric and ground field. Always in this framework a particular attention will be devoted to the investigation of the role of turbulence, intermittency and multifractal features of the geomagnetic disturbances and in situ satellite measurements (FP7 STORM project);
- The analysis of ULF waves (10-100 mHz) and geomagnetic pulsations in the solar wind, magnetosheath, magnetosphere, and at low and Antarctic latitudes. These studies will concentrate on the short and long term variations of the magnetospheric activity, through the analysis of long series of ULF magnetic measurements and geomagnetic pulsations, to investigate the possible influence on the different layers of the Earth's atmosphere. This research will benefit of the data from the new geomagnetic station at Talos Dome;
- The remote sensing of the plasma mass density in the plasmasphere by the resonance frequency of geomagnetic field lines. This information, together with observations from a network of whistler receivers, will be used to create a dynamical model of the Earth's plasmasphere (FP7 PLASMON project);
- The study of plasma processes in the ionosphere conditioned to solar wind conditions. It is foreseen to install the second radar at Dome C, the Dome C radar North (Italian PNRA - project), which will fill existing gaps in latitudinal coverage of SuperDARN network in the Southern Hemisphere.
- The study of the turbulence features of the magnetic field in the framework of the magnetosphere-ionosphere coupling and as a function of the geomagnetic activity level.
- Investigation of the features of magnetic field fluctuations also in planetary magnetospheric environments beyond the Earth's one, such as Mercury.



**DIVISION IV:**  
**“Solar Wind and Interplanetary Magnetic Field” (Coord.: R. Bruno)**

**A) Institutions involved in research activity**

- INAF-OATO (Turin) – studies of the extended corona and the origin, acceleration and transient perturbations of the solar wind. PI-role in ESA/Solar Orbiter-METIS instrument
- INAF-OAA & Astronomy Dpt. Univ. of Florence – studies of explosive phenomena, solar surface magnetism, solar instrumentation, and theory of the generation and transfer of polarised radiation in the solar atmosphere and the solar corona.
- INAF-IAPS (Rome) – studies related to solar wind plasma, solar energetic particles and cosmic rays; design and realization of plasma sensors; management of IAPS Large Plasma Chamber for ionospheric payloads plasma environmental tests; CoPI-role in ESA/Solar Orbiter-SWA(Solar Wind Analyzer) instrument
- INAF-OAR (Rome) – studies of solar irradiance; involved in the Conceptual Design Study of the EST; continuous management of the PSPT telescope (RISE project) for acquisition of photometric images of the whole Solar disk.
- INGV (Rome) – Space-Weather studies with particular emphasis on ionospheric perturbations following the IMF-solar wind coupling with the Earth’s magnetosphere
- Physics Dpt., Univ. Roma Tor Vergata – studies of the dynamics of solar photosphere; involved in the Conceptual Design Study of the EST; leading role in a small space mission in response to ASI-Small Missions Program call; participation to space weather awareness programme.
- Physics Dpt., Univ. L’Aquila – studies related to the solar wind properties;
- Physics Dpt., Univ. Cosenza – theoretical studies on space plasma kinetic properties; solar wind data analysis and modeling;
- INAF-OACt & Physics Dpt., Univ. Catania – studies of solar active regions and eruptive phenomena with data analysis and modeling; involved in the Conceptual Design Study of the EST.

**B) Scientific Report**

**B1) Main Scientific Themes**

**• Solar Wind and MHD Turbulence in Space Plasmas**

One of the central scientific interests of Division IV is the study of the solar wind and, more in general, the problem of MHD turbulence in space plasmas. In particular, the study has been devoted to the following topics: i) Statistical investigation of the MHD invariants of magnetic fluctuations in the interplanetary space, particularly within flux ropes, ii) radial evolution of density fluctuations intermittency during the expansion of the solar wind in the inner heliosphere, iii) radial evolution of the spectral break location between fluid and kinetic scales, iv) flattening of the spectral slope at proton scales moving from fast to slow wind, v) characterization of fluctuations within the kinetic regime as a function of wind speed and radial expansion, vi) study of the intermittency at proton scales.

Other studies have been made on the possible role that turbulence and magnetic coherent structures advected by the wind have in the modulation of Solar Energetic Particles produced by impulsive events.

Finally, the role of turbulence in solar wind heating has been evaluated estimating the energy transferred across the inertial range of the spectrum of the fluctuations to be dissipated at ion and/or electron scales. In addition, theoretical speculations, based on the available data, have been proposed about the nature of the fluctuations within the kinetic range.

Other studies have been conducted on reconnection events in the solar wind, 1) model vs observations for turbulence transport in the solar wind, 2) study of twisted magnetic flux-tubes in the solar wind.

- **Participation in ESA-Solar Orbiter Mission**

Within the participation to SOLAR ORBITER-SWA, the involvement is both managerial and scientific.

INAF-IAPS is the Italian leading Institute, at Co-PI level, for the national participation in the Solar Wind Analyser (SWA) experiment on board Solar Orbiter. The project is now in phase C-D and INAF-IAPS is working on the provision of the common Data Processing Unit which will manage the whole plasma suite made of 4 sensors dedicated to protons, electrons and minor ions.

Moreover, the related scientific activity mainly focusses on the use of numerical simulations apt to produce the ions and electrons velocity distributions for different solar wind conditions. The main goal is that of setting up the best measurement strategy to be adopted by SWA consistently with the DPU computational performances and the available telemetry.

- **Solar Photospheric Dynamics**

Another important scientific themes is the dynamics of solar photosphere, particularly the convective regime present in the outer layers of the sun. This topic is tackled by means of experimental techniques, images analysis procedures and numerical simulations:

Experimental techniques: realization of solar image acquisition systems based on CCD and CMOS detectors for panoramic Fabry-Pérot based monochromators (IBIS@DST/NSO) and achievement of observational campaigns at various ground-based telescope often coordinate with spatial instrumentation.

Image analysis: realization of automatic procedure of image segmentation (dynamic and optimal threshold, Medial Axis Transform, skeleton, spatial operators) and investigation of statistical tools for pattern analysis.

Numerical simulation: realization of dynamical models of passive magnetic advection.

Particular interest has been dedicated to estimate the possible contribution of the jets X to the solar wind. Mass flux has been evaluated after an accurate determination of the physical parameters. This study showed that jets, in particular blowout jets, are not sufficient for the solar wind. Also the contribution coming from the plumes has been estimated and, also in this case, an upper limit contribution, possibly of only 60%, has been determined.

- **Solar Energetic Particles**

Efforts have been devoted to study the acceleration and propagation of solar energetic particles (SEPs) in the interplanetary space as well as their interaction with the Mercury's environment. Moreover, the general features of the cosmic ray modulation, such as most of the step-like decreases and the Gnevyshev Gap phenomenon, have been investigated through non stationary data processing methods (e.g., empirical mode decomposition and wavelet analyses) in association with the variability of solar activity and the interplanetary magnetic field. The same analyses have been applied to the photospheric magnetic field data to understand the spatio-temporal dynamics of the solar magnetic field.

- **Space Weather**

This research, mainly performed by means of GPS sampling measurements at 50 Hz, allows the study of the ionospheric irregularities causing scintillation on the trans-ionospheric signals transmitted by GNSS satellites. The experimental observations can provide information useful for purely scientific advancements and for improve the predicting capabilities for Space Weather purposes. As a matter of fact, the original development of scintillation models needs to characterize the ionosphere also as function of the interplay between IMF-solar wind and ionosphere. Another topic on which this Division is working on is about the understanding of

the long-term variation of the ionosphere: the investigation of upper atmosphere secular trends, eventually connected to anthropogenic effects (greenhouse effect) and/or natural causes (next excursion or inversion of the geomagnetic field).

### **B2) Main Research Projects/Programmes**

- SOLAR ORBITER: Supporto Scientifico per la Realizzazione degli Strumenti METIS e SWA\_DPU. The related ASI contract for phase A-B1 ended in September 2011. The new contract for phase B2-C1 started in May 2012 and will last 3 years.
- EST: European Solar Telescope Conceptual Design Study financed by the European Commission
- SPARC: Space Awareness for Critical Infrastructure: European Commission - Directorate-General Home Affairs
- BEPI-COLOMBO: BepiColombo (SIMBIOSYS, ISA, PHEBUS, MEA, MIXS, SIXS) – The related ASI contract for phase B2/C will end in June 2013
- GRAPE (GNSS Research and Application for Polar Environment) Expert Group funded by SCAR ([www.grape.scar.org](http://www.grape.scar.org))

### **B3) Funding Agencies**

- European Commission: FP7 and Directorate-General
- ASI: Italian Space Agency
- SCAR: Scientific Committee for Antarctic Research

### **B4) Organization of Conferences and Workshop of international relevance**

- EWASS2012 Symposium 8 - The Sun: new tools and ideas in observational solar astrophysics
- EWASS2012 Special Session 11 - From solar physics to astrophysics: the Sun as Rosetta stone for understanding astrophysical processes
- SOC-5th Earth-Sun-Exploration conference, hold in January 2013
- Co-Chair of Solar Wind XIII-2012-Session II – “The dynamical behavior of the interplanetary medium throughout the heliosphere, including large- and small-scale structures, the evolution of disturbances, dissipative processes, including turbulence, and energetic particles”;
- Chair of Solar Wind XIV-2015-Session II “Solar wind (quiescent and transient) evolution: processes, structures, waves, and turbulence”
- ISSS course on: “Frontiers of Space Science: from Solar Activity to NEOs” held in L’Aquila, (17 – 22 April 2011) directed by F. Berrilli, A. Celletti (University of Roma “Tor Vergata”), E. Flamini (ASI) and D. Mueller (ESA).
- ISSS course on: “Astrophysical and Space Plasmas” held in L’Aquila, (2 – 7 September 2012) directed by A. Ferrari (CIFS, Torino & University of Torino), M. Tavani (INAF-IAPS & University of Roma “Tor Vergata”), B. Coppi (M.I.T., Cambridge , USA), R. Rosner (University of Chicago, USA)
- ISSS Course on: “Towards a Deeper Understanding of the Sun and the Heliosphere with Solar Orbiter – The first Solar Orbiter School” directed by R. Bruno (INAF-IAPS), E. Antonucci (INAF-OATO) and D. Mueller (ESA), L’Aquila, 22-25 Sept. 2014
- Science Working Team of the Solar Orbiter mission organized by European Space Agency, L’Aquila 25-27 Sept. 2014

### **B5) Prizes/Recognitions**

- D. Telloni (INAF-OATO) received the Second Edition (2011) of Premio Ferraro Award, for the best PHD thesis on MHD studies in astrophysical or laboratory plasmas
- The Minor Planet Center (MPC) under the auspices of the IAU Division III designates the asteroid 7918 = 1981 EJ22 with the name Berrilli.

### **C) Goals, priorities and plans for future activities**

- **SOLAR ORBITER:** the ASI scientific contract “Supporto Scientifico per la Realizzazione degli Strumenti METIS e SWA\_DPU nelle fasi B2-C1” started in May 2012 and will end in May 2015. A new contract is expected to start afterwards to last another three years, covering the delivery phase of SWA FM to ESA. In the meantime, the scientific team supports the industry during the production of the required deliverables.
- **BEPI-COLOMBO:** recently, because of financial difficulties, ASI decided not to support further activities related to Bepi-Colombo not having a PIship or CoPIship involvement in this program. Despite that, the scientific collaboration on energetic particles will continue as much as possible on the simulation of the propagation of SEPs in Mercury’s magnetosphere as well as of the flux of secondary particles produced by interaction with the planetary surface. In addition, evolution of SEP events of different intensities will be studied for the correlation between the observations of the SEP radiation monitor, designed for the LISA Pathfinder mission, and the particle fluxes charging the gravitational sensor test masses.
- **STORM(FP7):** Solar system plasma Turbulence: Observations, intermittency and Multifractals. This project will use large datasets instead of focusing on particular events, to advance the understanding of fundamental processes such as the turbulent energy transfer, the intermittent turbulence and multifractals in astrophysical plasmas.
- **SOLARNET(FP7):** High Resolution Solar Physics Network: is the solar successor of OPTICON, and start in April 2013. Main objectives are to provide access to European Solar telescopes and instruments, and to develop data pipelines and innovative instrumentations such that science-ready data is easily available for users of the ACCESS program.
- **MISW(FP7):** Mitigation of space weather threats to GNSS services (2013-2015)

## DIVISION V:

### “Geomagnetic Observatories, Surveys and Analyses” (Coord.: A. Meloni)

#### 1) GEOMAGNETIC OBSERVATORIES

In Italy, for the years of concern, INGV has taken care of activities of 6 magnetic observatories:

| Observatory                | IAGA code | Latitude | Longitude | Elev. (asl m) |
|----------------------------|-----------|----------|-----------|---------------|
| <u>L'Aquila</u>            | AQU       | 42°23'N  | 13°19'E   | 682           |
| <u>Castello Tesino</u>     | CTS       | 46°03'N  | 11°39'E   | 1175          |
| <u>Mario Zucchelli</u> (*) | TNB       | 74°42'S  | 164°6'E   | 30            |
| <u>Concordia</u> (**)      | DMC       | 75°06'S  | 123°21'E  | 3200          |
| <u>Duronia</u>             | DUR       | 41°39'N  | 14°28'E   | 918           |
| Lampedusa                  | LMP       | 35°31'N  | 12°32'E   | 33            |

All observatories are operated by INGV except (\*) in cooperation with PNRA and (\*\*) in cooperation with PNRA and France, IPEV.

#### • L'AQUILA

On April 6th 2009, L'Aquila and surroundings, were struck by a very bad M=6.2 earthquake that destroyed the city and killed 309 people. This Observatory, located near Preturo, ten km north-west from the city, was not severely affected by earthquake but the land on which the observatory buildings: a) absolute measurements, b) variometers, c) proton precession vector magnetometer, d) laboratory and e) general services, were built, was asked back by L'Aquila University (the original landlord) for new student housings. L'Aquila was an INTERMAGNET Observatory and for this observatory, K magnetic activity indices, SSC, SFE and SI notifications were made. As now we are still waiting for an official request for moving out, but standard INTERMAGNET activity was interrupted.

The Observatory started its activity in 1958 and yearbooks from 1960 onwards were published; last published as now is 2009. Today the main observatory in central Italy is located in Duronia, but a complete system is still running in the buildings of this old geomagnetic observatory. Observations are made by means of an Overhauser magnetometer for the measurement of the field intensity F and a fluxgate magnetometer, for the measurement of H, D and Z component variations. The instruments have a 0.1 nT resolution; original sampling rate is 1 Hz, then the data are filtered with a Gauss filter and recorded at 1 min.

#### • CASTELLO TESINO

The Observatory of Castello Tesino (Northern Italy) is located about fifty-five km East from Trento. It has been working almost continuously since 1964. It works now as an automatic Observatory and consists of three buildings, completely non-magnetic; a) laboratory for small repairs, b) absolute measurement equipments, and c) the automatic digital variometer system. Since some disturbance affected the Observatory variometer house, recently a new small wooden building was built at a safe distance, in order to avoid this problem. After a trial period, the old house was finally left in April 2012. In the same period the power supply was converter from AC 220 V to DC 12 V in order to improve the quality of measurement systems. Two systems are currently working simultaneously in the new wooden building.

Declination and Inclination measurements are carried out by means of an optical theodolite equipped with a fluxgate magnetometer. The horizontal and vertical intensities are determined associating the I value to the total field measurements made by the scalar magnetometer. The geomagnetic field is recorded by means of two parallel systems, each constituted by a proton precession magnetometer (Overhauser GEM-GSM90) for the measurement of the total magnetic field intensity F and a tri-axial fluxgate magnetometer (LEMI-017) for the measurement of the H, D and Z component variations. The instruments have a 0.1 nT resolution; the original sampling rate is 1 Hz, then the data are filtered with a Gauss filter and recorded at 1 min. At this time all yearbooks from 1996 to 2013 were completed, all regularly published as standard booklets. From 2010 yearbooks are not printed but are available on the website of INGV.

At this time the yearbooks from 1996 to 2013 are available on the INGV website ([http://roma2.rm.ingv.it/en/facilities/geomagnetic\\_observatories](http://roma2.rm.ingv.it/en/facilities/geomagnetic_observatories)) where the monthly bulletins are also available.

- **TERRA NOVA BAY (MARIO ZUCHELLI STATION)**

This Observatory was installed during 1986-87 austral summer at the Italian Antarctic Mario Zucchelli Station. In the first years geomagnetic field measurements were carried out only during summer expeditions. Since 1991 the recording was implemented with an automatic acquisition system operating through the year. Declination and Inclination measurements are carried out by means of an optical theodolite equipped with a fluxgate magnetometer. The horizontal and vertical components are determined associating the I value to the total field measurement made by a proton precession magnetometer. A proton precession magnetometer for the measurement of F and a fluxgate magnetometer for the measurement of the H, D and Z component variations, are used for time variations; both instruments have a 0.1 nT resolution; the original sampling rate is 1 Hz, then the data are filtered with a Gauss filter and recorded at 1 min.

When absolute measurements are available (summer) results, including hourly mean data, are published on pdf booklets (so far completed up to 2010-2011). Results are available at INGV website:  
[http://roma2.rm.ingv.it/it/risorse/banche\\_dati/32/dati\\_osservatorio\\_di\\_stazione\\_mario\\_zucchelli/46/annuari](http://roma2.rm.ingv.it/it/risorse/banche_dati/32/dati_osservatorio_di_stazione_mario_zucchelli/46/annuari).

- **CONCORDIA (CONCORDIA STATION)**

In 1994 France and Italy started a program for opening a permanent scientific station on the high East Antarctic plateau at Dome C at 3280 m asl. The national Antarctic Programs (IPEV and PNRA respectively) started logistic, technical and scientific activities at Dome C that were initiated with the realization of a summer camp. The permanent station opened in 2005.

The observatory is constituted by two shelters, a) variometer shelter and b) absolute measurements shelter, at a distance of about 300 m from the old field camp. Operations started regularly at the end of 2004. Variations of the Earth's magnetic field are monitored by a three-axis fluxgate magnetometer along three orthogonal vector components oriented with respect to the local magnetic meridian. The intensity of the field is measured by an Overhauser magnetometer. Absolute measurements are performed during the whole year.

The flux-gate variometer, a suspended DMI magnetometer, and an Overhauser magnetometer operate for the acquisition of the geomagnetic field intensity and components time variations. The instrument sensors are located in a cave under the shelter in order to keep the sensors at a constant temperature (about -40°C). In the absolute shelter D, I, F absolute measurements are carried out according to standard observatory practice. Data quality and standard requirements have allowed the inclusion of Concordia as an INTERMAGNET Observatory.

- **DURONIA**

Since the end of 2007 an electromagnetic field monitoring station has been in operation at Duronia (Lat. 41°39'N, Lon. 14°28'E). The station was created in the framework of a project headed by the Abruzzo region. The main target was to create a network of stations to monitor the environmental electromagnetic signals in the Adriatic area, in the frequency band from 0.001Hz to 100kHz (ULF-ELF-VLF). The peculiarity of Duronia installation is the site low electromagnetic background noise and the low noise of the instrumentation. A target will also be the long-term monitoring of local magnetic field anomalies possibly related to the local geodynamical processes. After the 2009 L'Aquila earthquake, Duronia station was completed to include a full geomagnetic equipment to follow IAGA requirements. On August 28, 2012 Duronia has been accepted as an INTERMAGNET Magnetic Observatory (IMO).

- **LAMPEDUSA**

By the end of 2007 magnetic field time variations are also recorded in Lampedusa Island South-West of Sicily (Lat. 35° 31' N, Lon. 12° 32' E). From the beginning of 2007 values of the Earth's magnetic field were recorded quite regularly. A small stone building with a wooden roof, located within a natural reserve, in an area characterized by a low electromagnetic noise, host the magnetic instruments. Electric current is provided by photovoltaic cells and data transmission is achieved through GSM connection. The little hut hosts only the instruments electronic units; sensors are buried in thermally isolated shafts in the area in front of the building. Total intensity F is measured by an Overhauser magnetometer, while the variations of the magnetic field components H, D and Z are measured by two fluxgate vector magnetometers. Data acquisition is made through a device made on purpose at INGV to avoid the use of a personal computer. Data can be found in the INGV website ([http://roma2.rm.ingv.it/it/risorse/osservatori\\_geomagnetici](http://roma2.rm.ingv.it/it/risorse/osservatori_geomagnetici)) where the monthly bulletins are also available.

## 2) GEOMAGNETIC STATIONS

Three geomagnetic time varying stations were in operation in Italy in the time period.

A geomagnetic station was installed in Cittareale (Lat. 42.62° N - Long. 13.16° E , 960 m asl) on 23 March 2011 and from that time has been recording continuously three components plus scalar intensity of the geomagnetic field. Its objective is not only just geomagnetic but also to eventually detect any anomaly associated with earthquakes, the place of installation being a seismic area with some frequent seismicity.

Another magnetometric system (formed by a couple of vectorial and scalar magnetometers) is running aboard of the seafloor GEOSTAR-class cabled Observatory (called NEMO-SN-1) which was deployed on 9 June 2012, 25 km offshore Catania (Sicily) (Lat. 37.54765°N, Lon. 15.3975°E, 2010 m depth). Data are streaming on the web through Catania harbor centre of INFN (National Institute of Nuclear Physics). The objective of the magnetic instruments is twofold: a) the magnetic monitoring of the seafloor environment in order to extend the national land network to the sea, b) to assist the other geophysical and oceanographic monitoring in order to check if there are magnetic anomalies eventually associated to extreme events typical of this area such as Etna volcanic eruptions and earthquakes.

Recently, a new multiparametric station including magnetic measurements was set up in Varese Ligure (La Spezia). Currently, the magnetic measurements have been recently analyzed and the results can be considered acceptable.

## 3) MAGNETIC SURVEYS

The spatial variation of the geomagnetic field and details on local secular variation over Italy, is achieved by means of a national magnetic network of measuring points. This

complements the observatory network. The Italian repeat station network consists of more than 110 points. INGV has the task to make measurements and data elaboration. The measurements are generally carried out by means of Declination/Inclination theodolites and total intensity magnetometers; a gyroscope theodolite is used to check and establish new azimuth marks when necessary. The information on the more rapid time variations, both for the diurnal variation and for possible irregular perturbations, is taken at the Italian geomagnetic observatories (Duronio, Castello Tesino and Lampedusa). Moreover, for selected areas, other temporary magnetic time recording stations, properly displaced to have nearly a total coverage of the Italian territory, are installed during the survey. The last full survey was made for the 2010.0 date. It was completed in November 2010 and consisted of 131 repeat stations, with 55 km average spacing, over Italy, and included also stations surveyed in Albania, Malta and Corsica. After data elaboration was completed a 1/2000,00 cartography for F, H, Z and D was realized. All maps are now published together with a CD rom and interactive program to display maps and magnetic field values across all Italy. A partial survey was made for the 2012.5 date. This most recent partially survey, that was completed in December 2012, consisted of 25 repeat stations and D, I and F magnetic element values on all stations have been obtained. Data compared with the 1/2000,00 cartography for F, H, Z and D are published together with a CD rom and interactive program to display maps and magnetic field values across all Italy. Now INGV is working at the measurements for the 2015.0 magnetic network, with field activities on repeat stations. This phase will be completed in 2015 and will lead to the 2015.0 magnetic elements absolute values, secular variation models and updated maps.

#### **4) OTHER ACTIVITIES RELEVANT TO DIVISION V**

**Geomagnetic Regional Reference Models:** As well known magnetic modeling provides scientific communities with updated information on magnetic field elements as the output of computer programs. They are based on updated measurements taken in the region under modelling. An analytical expression, a second order polynomial, in latitude and longitude for the field elements was determined and coefficients for 2010.0 with average secular variation over the period 2005-2010, were obtained for Italy. Moreover the new maps for Italy, for D, F, H and Z at the epoch 2010.0, were also produced. ARM (Antarctic Reference Model) is a 3D model for the geomagnetic field over Antarctic regions. It is based on the Spherical Cap Harmonic Analysis and allows the computation of the main field and its secular variation over these regions from 1960 onwards. Another regional model with similar characteristics of ARM has been developed to model the geomagnetic field of Southern Italy and Albania (including the confining sea) since 1990. Both models ARM and this latter model have been now updated to include year 2010 with prediction to 2012.

**Crustal magnetic field investigations.** The study of tectonics can take important advantages by the use of magnetic field surveys that allow the determination of the crustal field contribution. We report here some recent contributions in this field. High spatial resolution aeromagnetic surveys were undertaken, especially at INGV, by the use of helicopter born magnetometry and data interpretation in volcanic magnetized areas. During the recent cruise MAVA2011, performed on the R/V Urania, a new magnetic dataset was obtained from seamounts Vavilov and Marsili. At the Eolie islands archipelago, in the Tyrrhenian, surveys and studies were made on Salina island and Volcano island. In the Tyrrhenian Sea, in the marine facing on Panarea and Stromboli islands, an high resolution magnetic survey (PANSTR12) was carried out using an Italian Navy ship and marine magnetometers with Overhauser effect sensors.

A group at the Università of Camerino has undertaken analysis and interpretation of marine and in particular devoted to the realization of kinematic models for the Mediterranean and the Atlantic.



## **“Interdivisional Commissions on History and on Education and Outreach” (Coord.s.: A. De Santis, F. Berrilli)**

As mentioned in the IAGA website, the Interdivisional Commission on History encourages historical research by scientists as well as professional historians of science - into the history of geophysics. It encourages historical sessions and presentations at IAGA meetings, and the preservation of IAGA's history. The Interdivisional Commission on Education and Outreach has become an autonomous interdivisional commission from the former from IAGA 2013 Assembly in Merida (Mexico). In this report they are considered merged together because they were in this way till 2013.

### **A) Institutions involved in research activity**

- Istituto Nazionale di Geofisica e Vulcanologia (INGV)
- Consiglio Nazionale delle Ricerche (CNR)
- ISPRA
- University of Tor Vergata
- Department of Civil Protection (DPC)

### **B) Activity Report**

#### **B1) Main Themes**

- INGV Open day (“Open Science for Open minds”) 21 May 2011. Exhibits on Geophysics, Geomagnetism and Paleomagnetism.
- SCIENZA APERTA (“Open Science”) 2012 -2014 (all INGV centers). In particular: different Exhibits on Geomagnetism and Paleomagnetism (INGV, Rome) and Geomagnetic Prospection (INGV, Catania)
- SHINE! (Scientist are Humans - Interactive Night of Entertainment) Sept. 2012-2014. Outreach Events, seminars, museums, exhibits in 24 Italian cities. INGV participated with initiatives in all INGV centers.
- Realization of Podcast: “Stromboli: a small volcano but a great natural laboratory” by INGV – Catania. This Podcast shows the magnetic monitoring of Stromboli volcano. 2012.
- Series of seminars given by INGV researchers (Dep. Rome 2) on Space Weather, Ionospheric Physics at the University La Sapienza, Rome Dec. 2012- Jan. 2013.
- Visits of school pupils/students/teachers in 2012-2014 at INGV: around 5000 people every year visited seven INGV centers, among which the main were Rome, Naples and Catania Centers.
- During the outreach event "Under another optics" the telescope in carbon fiber built during the Stage in heliophysics at Tor Vergata was presented April 6, 2013, in the High School of Ceccano (FR).
- Laboratory of Solar Physics at University of Tor Vergata Stages 2011, 2012, 2013, 2014, for teachers and students from secondary schools. The laboratory section of the stage was executed in two phases (weeks): i) First phase aims was the realization of a keplerian telescope and low-cost acquisition system. During this week students were introduced to astronomical techniques used to safely collect and acquire solar light; ii) Second phase aims was the realization of a low-cost instrument to analyse sunlight extracting information about the solar spectrum, solar irradiance and Sun. <http://www.stageatorvergata.it/home>

#### **B3) Funding Agencies**

- University and Research Ministry (MIUR)
- Italian National Programme for Antarctica Research (PNRA)

- Italian Space Agency (ASI)
- European Commission (FP7-Space Call)
- Piano Lauree Scientifiche (PLS)

#### **B5) Prizes**

- Edmund Optics (EO), a provider of optical components for industry and academia, has announced the winners of its 2013 Higher Education Global Grant Program. The awards are given in recognition of outstanding undergraduate and graduate optics programs in science, technology, engineering, and mathematics at non-profit colleges and universities worldwide. Third-place European recipient is Università degli Studi di Roma "Tor Vergata" (Italy), submitted by Luca Giovannelli. The award was presented for developing a Fabry-Pérot Capacitance Stabilized Etalon (CSE), based on a monolithic approach, optimized for a satellite application.

#### **C) Goals, priorities and plans for future activities**

- Promotion and Initiatives on History of Geomagnetism and Aeronomy, as well as in Education and Outreach will continue during the next two years, repeating in particular the positive experience of SCIENZA APERTA ("Open Science") and SHINE!. Some of the main centers of INGV will be again open to school visits, especially in Rome with three visits/week from October of each year to June of the next.
- Laboratory of Solar Physics at University of Tor Vergata Stages 2015-2016 for teachers and students from secondary schools. For these laboratories is planned the realization of solar radiotelescopes for education and connection with solar activity. More detailed and updated information can be found on the web (e.g. <http://www.stageatorvergata.it/home> or <https://www.fisica.uniroma2.it/~solare/english/>).

## SCIENTIFIC PUBLICATIONS

### DIVISION I:

#### “Internal Magnetic Field”

##### 2011

1. Agnini C., E. Fornaciari, L. Giusberti, P. Grandesso, L. Lanci, V. Luciani, G. Muttoni, D. Rio, C. Stefani, H. PŠlike and D.J.A. Spofforth, (2011) Integrated bio-magnetostratigraphy of the Alano section (NE Italy): a proposal for defining the Middle-Late Eocene boundary, *Geol. Soc. Am. Bull.* 123, 5/6, 841–872; doi: 10.1130/B30158.1.
2. Balasco M., Galli P., Giocoli A., Gueguen E., Lapenna V., Perrone A., Piscitelli S., Rizzo E., Romano G., Siniscalchi A., Votta M. (2011). Deep geophysical electromagnetic section across the middle Aterno Valley (central Italy): preliminary results after the April 6, 2009 L’Aquila earthquake. *Bollettino Di Geofisica Teorica e Applicata*, vol. 52(3), p. 443-455, ISSN: 0006-6729, doi: 10.4430/bgta0028
3. Benhamou S., Sudre J., Bourjea J., Ciccione S., De Santis A., and Luschi P., The role of geomagnetic cues in green turtle open sea navigation, *PLoS One*, vol. 6, N.10, 1-11, 2011.
4. Barraclough D.R., De Santis A., (2011) Repeat stations activities, in *Geomagnetic Observations and Models*, M. Manda and M. Korte (Ed.s), IAGA Special Sopron Book Series, vol. 5, 45-55, Springer.
5. Boncio P., A. Pizzi, G. Cavuoto, M. Mancini, T. Piacentini, E. Miccadei, G.P. Cavinato, S. Piscitelli, A. Giocoli, G. Ferretti, R. De Ferrari, M.R. Gallipoli, M. Mucciarelli, V. Di Fiore, F. Pergalani, G. Naso and Working Group Macroarea3, B. Giaccio, M. Moscatelli, M. Spadoni, G. Romano, A. Franceschini, D. Spallarossa, M. Pasta, M. Pavan, D. Scafidi, S. Barani, C. Eva, M. Compagno, T. Campea, G.R. Di Bernardino, T. Mancini, A. Marino, R. Montefalcone, F. Mosca. Geological and geophysical characterisation of the Paganica – San Gregorio area after the April 6, 2009 L’Aquila earthquake (Mw 6.3, central Italy): implications for site response. *Boll. Geof. Teor. e Appl.* Vol. 52 (2011), n. 3, pp. 491-512;
6. Capraro L., Massari, F., Rio, D., Fornaciari, E., Backman J., Channell J.E.T., Macrì, P., Prosser, G. and Speranza, F. (2011), Chronology of the Lower-Middle Pleistocene succession of the south-western part of the Crotone Basin (Calabria, Southern Italy). *Quat. Sci. Rev.*, 30, 1185-1200, doi: 10.1016/j.quascirev.2011.02.008.
7. Cicchino A., Zanella E., De Astis G., Lanza R, Lucchi F., Tranne C.A., Airoidi, G., Mana S., (2011). Rock magnetism and compositional investigation of Brown Tuffs deposits at Lipari and Vulcano (Aeolian Islands–Italy). *Journal of Volcanology and Geothermal Research*, DOI 10.1016/j.volgeores.2011.08.007
8. Cifelli F., Minelli L., Rossetti F. & Mattei M., 2011. The emplacement of the Late Miocene Monte Capanne intrusion (Elba Island, Central Italy): constraint from magnetic fabric analyses. *Int. J. Earth Sci.*, doi 10.1007/s00531-011-0701-z.
9. Cosentino, D., Bertini, A., Cipollari, P., Florindo, F., Gliozzi, E., Grossi, F., Lo Mastro S., and Sprovieri M. (2011). Orbitally-forced palaeoenvironmental and palaeoclimate changes in the late post-evaporitic Messinian stage of the central Mediterranean Basin. *Geol. Soc, Am, Bull*, 2011B30462.1, doi: 10.1130/B30462.1
10. D'Ajello Caracciolo, F., A. Pignatelli, F. Speranza and A. Meloni (2011), A re-evaluation of the Italian historical geomagnetic catalogue: implications for paleomagnetic dating at active Italian volcanoes. *Solid Earth*, 2, 65-74.
11. Dela Pierre, E. Bernardi, S. Cavagna, P. Clari, R. Gennari, A. Irace, F. Lozar, S. Lugli, V. Manzi, M. Natalicchio, M. Roveri, D. Violanti (2011). The record of the Messinian salinity crisis in the Tertiary Piedmont Basin (NW Italy): The Alba section revisited. *Paleogeogr. Paleoclimatol. Palaeocol.*, vol. 310, p. 238-255, ISSN: 0031-0182
12. De Santis A., Qamili E., Cianchini G. (2011) Ergodicity of the recent geomagnetic field, *Phys. Earth*

13. Di Mauro, D., L. Alfonsi, V. Sapia, L. Nigro and M. Marchetti (2011), First Field Magnetometer Investigation at the Phoenician Island of Mozia (Trapani), Northwestern Sicily: Preliminary Results, *Archaeological Prospection*, Volume 18, Issue 3, pages 215–222, DOI: 10.1002/arp.41
14. D’Orlando F., L. Angeletti, L. Capotondi, M. A. Laurenzi, M. Taviani, L. Torelli, T. Trua, L. Vigliotti and N. Zitellini. (2011). Coral Patch and Ormonde seamounts as a product of the Madeira hotspot, Eastern Atlantic Ocean. *Terra Nova*, Vol. 22 (6), 494-500.
15. Galeotti, S., Lanci, L., Florindo, F., Naish, T., Sagnotti, L., Sandroni, S., Talarico, F. (2011). Cyclochronology of the Eocene-Oligocene transition from a glacial-marine succession off the Victoria Land coast, Cape Roberts Project, Antarctica. *Palaeogeogr., Palaeoclimatol., Palaeocol.*, doi:10.1016/j.palaeo.2011.08.011.
16. Gioia D., Schiattarella M., Mattei M. & Nico G. (2011). Quantitative morphotectonics of the Pliocene to Quaternary Auletta basin, southern Italy. *Geomorphology*, 134, 3-4, 26–34, 3.doi:10.1016/j.geomorph.2011.07.009.
17. Gurioli L., Zanella E., Gioncada A., Sbrana A. The historic eruption of “Breccia di Commenda” (Vulcano, Italy). *Bulletin of Volcanology*.
18. Iaccarino S. M., A. Di Stefano, L. M. Foresi, E. Turco, N. Baldassini, A. Cascella, S. Da Prato, L. Ferraro, R. Gennari, F. J. Hilgen, F. Lirer, R. Maniscalco, R. Mazzei, F. Riforgiato, B. Russo, L. Sagnotti, G. Salvatorini, F. Speranza, M. Verducci (2011) High-resolution integrated stratigraphy of the upper Burdigalian-lower Langhian in the Mediterranean: the Langhian historical stratotype and new candidate sections for defining its GSSP. *Stratigraphy*. vol. 8, nos. 2-3, pp. 199-215.
19. Kirscher U., Aubele K., Muttoni G., Ronchi A. and Bachtadse V. (2011). Paleomagnetism of Jurassic Carbonate Rocks from Sardinia – No Indication of Post Jurassic Internal Block Rotations. *Journal of Geophysical Research*. Vol. 116, B12107, Doi:10.1029/2011jb008422.
20. Lanci L., Delmonte B., Maggi V., Kent D.V., Biscaye P.E., Petit J-R. (2011). Magnetization of polar ice: a measurement of terrestrial dust and extraterrestrial fallout, *Quaternary Sci. Rev.*
21. Ledo J., Jones A. G., Siniscalchi A., Campaña J., Kiyani D., Romano G., Rouai M. and TOPOMED MT TEAM (Arbolea, Balasco, Guimera, Hogg, Lapenna, Marcuello, Moretti, Pina, Queralt, Rosell, Seddas, Schmoldt, Teixell, Tripaldi) (2011). Electrical signature of modern and ancient tectonic processes in the crust of the Atlas mountains of Morocco. *Physics of The Earth and Planetary Interiors*, vol. 185, p. 82-88, ISSN: 0031-9201, doi: 10.1016/j.pepi.2011.01.008.
22. Lesti, C., Porreca, M., Giordano, G., Mattei, M., Cas, R., Wright, H., Viramonte J. (2011) - High temperature emplacement of the Cerro Galán and Toconquis Group ignimbrites (Puna plateau, NW Argentina) determined by TRM analyses. *Bulletin of Volcanology*. 73,1535-565. DOI 10.1007/s00445-011-0536-2.
23. Molina, E., Alegret, L., Apellaniz, E., Bernaola, G., Caballero, F., Dinarès-Turell, J., Hardenbol, J., Heilmann-Clausen, C., Larrasoña, J.C., Luterbacher, H., Monechi, M., Ortiz, S., Orue-Etxebarria, X., Payros, A., Pujalte, V., Rodríguez-Tovar, F.J., Tori, F. and Tosquella, J. (2011), The Global Standard Stratotype-section and Point (GSSP) for the base of the Lutetian Stage at the Gorrondatxe section (Spain). *Episodes*, 32, 86-108.
24. Muttoni G., G. Scardia, D.V. Kent, E. Morsiani, F. Tremolada, M. Cremaschi, C. Peretto. First dated human occupation of Italy at ~0.85 Ma during the late Early Pleistocene climate transition. (2011). *Earth and Planet. Sci. Lett.* 307, 241–252.
25. Nomade, S., Muttoni, G., Guillou, H., Robin, E., Scardia, G. (2011). First <sup>40</sup>Ar/<sup>39</sup>Ar age of the Ceprano man (central Italy). *Quaternary Geochronology*, 6, 453–457.
26. Omodeo Salè S., Gennari R., Lugli S., Manzi V., Roveri M.. Tectonic and climatic control on the Late Messinian sedimentary evolution of the Nijar Basin (Betic Cordillera, Southern Spain): implications for the salinity crisis in the Mediterranean. *Basin Research*
27. Pavia M., Zunino M., Coltorti M., Angelone C., Arzarello M., Bagnus C., Bellucci L., Colombero S., Marcolini F., Peretto C., Petronio C., Petrucci M., Pieruccini P., Sardella R., Tema E., Villier B., Pavia

- G. (2011). Stratigraphical and paleontological data from the Early Pleistocene Pirro 10 site of Pirro Nord (Puglia, south-eastern Italy). *Quaternary International*, 1-16, doi: 10.1016/j.quaint.2010.12.019.
28. Payros, A., Dinarès-Turell, J., Bernaola, G., Orue-Etxebarria, X., Apellaniz, E. and Tosquella, J. (2011), On the age of the Early/Middle Eocene boundary and other related events: cyclostratigraphic refinements from the Pyrenean Otsakar section and the Lutetian GSSP. *Geol. Mag.*, 148, 442-460, doi: 10.1017/S0016756810000890.
  29. Roberts, A.P., Chang, L., Rowan, C., Horng, C.-S., Florindo, F., (2011). Magnetic properties of sedimentary greigite (Fe<sub>3</sub>S<sub>4</sub>): An update. *Reviews of Geophysics*, 49, RG1002, doi: 10.1029/2010RG000336
  30. Roberts, A.P., Florindo, F., Giuliana Villa, Liao Chang, Luigi Jovane, Steven M. Bohaty, Juan C. Larrasoana, and John D. Fitz Gerald (2011), Magnetotactic bacterial abundance in pelagic marine environments is limited by availability of dissolved iron and organic carbon flux, *Earth Planet. Sci. Lett.*, 310, 441-445.
  31. Sagnotti L. (2011), Magnetic Anisotropy, in: Encyclopedia of Solid Earth Geophysics, (Editor Harsh K. Gupta), Springer, 1540 pp., p. 717-728.
  32. Sagnotti L., Smedile A., De Martini P.M., Pantosti D., Speranza F., Winkler A., Del Carlo P., Bellucci L.G. and Gasperini L. (2011), A continuous paleosecular variation record of the last 4 millennia from the Augusta Bay (Sicily, Italy). *Geophysical Journal International*, 184, 191-192, doi:10.1111/j.1365-246X.2010.04860.
  33. Sagnotti, L., Macrì P., Lucchi, R., Rebesco, M., and Camerlenghi A. (2011), A Holocene paleosecular variation record from the northwestern Barents Sea continental margin, *Geochem. Geophys. Geosyst.*, 12, Q11Z33, doi:10.1029/2011GC003810.
  34. Saragnese F., L. Lanci and R. Lanza, (2011) Nanometric-sized atmospheric particulate studied by magnetic analysis, *Atmospheric Environment*, 45, 450-459.
  35. Speranza, F., P. Macrì, D. Rio, E. Fornaciari, and C. Consolaro (2011), Paleomagnetic evidence for a post-1.2 Ma disruption of the Calabria terrane: consequences of slab break-off on orogenic wedge tectonics. *Geol. Soc. Am. Bull.* 123, 925-933, doi:10.1130/B30214.1.
  36. Tema E. & Kondopoulou D. (2011). Secular variation of the Earth's magnetic field in the Balkan region during the last eight millennia based on archaeomagnetic data. *Geophysical Journal International*, 186, 2, pp.603-614, doi: 10.1111/j.1365246X.2011.05088.x.
  37. Tema E. (2011). Archaeomagnetic Research in Italy: Recent achievements and future perspectives. In: The Earth's Magnetic Interior, IAGA Special Sopron Book Series, Volume 1, Chapter 15, pp. 213-233. Eds: Petrovsky, E., Herrero-Bervera, E., Harinarayana, T., Ivers, D., Springer, doi: 10.1007/978-94-007-0323-0-15.
  38. Turco E., Cascella A., Gennari R., Hilgen F.J., Iaccarino S.M. and Sagnotti L. (2011), Integrated stratigraphy of the La Vedova section (Conero Riviera, Italy) and implications for the Burdigalian/Langhian boundary, *Stratigraphy*, vol. 8, nos. 2-3, pp. 89-110.
  39. Venuti, A., Florindo, F., Caburlotto, A., Strada, E., Talarico, F., Hillenbrand, C.-D., Hounslow, M. (2011), Late Quaternary sediments from deep-sea sediment drifts on the Antarctic Peninsula Pacific margin: Climatic control on provenance of minerals, *J. Geophys. Res.*, 116, B06104, doi:10.1029/2010JB007952.
  40. Vigliotti, L., A. Ascoli, C. Bergami, L. Capotondi & A. Piva. (2011). Magnetic properties of the youngest sapropel S1 in the Ionian and Adriatic Sea: inference for the timing and mechanism of sapropel formation. *Ital.J.Geosci.*, Vol. 130, No. 1. 106-118. DOI: 10.3301/IJG.2010.29
  41. Zanella E., Cicchino A., Lanza R. (2011) Composite detrital and thermal remanent magnetization in tuffs from Aeolian Islands (southern Tyrrhenian Sea) revealed by magnetic anisotropy. *International Journal of Earth Sciences*, DOI 10.1007/s00531-011-0651-5
  42. Zanella E., Ferrara E., Bagnasco L., Ollà A., Lanza R., Beatrice C. (2011). Magnetite grain-size analysis and sourcing of Mediterranean obsidians. *Journal of Archaeological Science*.

## 2012

43. Airoidi G., Muirhead J.D., Zanella E., White J.D.L. (2012) Emplacement process of Ferrar Dolerite sheets at Allan Hills (South Victoria Land, Antarctica) inferred from magnetic fabric. *Geophys. Jour. Int.*, Vol. 188, Issue 3, 1046–1060.
44. Alfonsi L., F.R. Cinti, D. Di Mauro, S. Marco (2012), Archaeoseismic Evidence of Two Neolithic (7,500–6,000 B.C.) Earthquakes at Tell es-Sultan, Ancient Jericho, Dead Sea Fault. *Seismological Research Letters*, 83, 4, 639-648.
45. Cascella A., Lirer F., Cesarano M., Casciello E. & Pappone G. (2012). Integrated stratigraphy of middle-late Miocene synorogenic deposits of the southern Apennines chain: the San Bartolomeo Flysch. *Riv. It. Paleont. Strat.* 118, 2, 295-311.
46. Cianchini G., De Santis A., D. R. Barraclough, L. X. Wu, and K. Qin, (2012). Magnetic transfer function entropy and the 2009 Mw = 6.3 L'Aquila earthquake (Central Italy), *Nonlin. Processes Geophys.* 19, pp. 401-409, doi:10.5194/npg-19-401-2012
47. Cifelli, F., Minelli, L., Rossetti, F. & Mattei, M. (2012) - The emplacement of the late Miocene Monte Capanne intrusion (Elba island, Central Italy): constraint from magnetic fabric analyses. *International Journal of Earth Sciences*. 101, 863–877, DOI: 10.1007/s00531-011-0701-z.
48. De Santis A., E. Qamili, G. Spada, P. Gasperini, (2012). Geomagnetic South Atlantic Anomaly and global sea level rise: a direct connection?, *J. Atm. Solar-Terr. Phys.*, 74, 129-135.
49. Di Chiara, A., Speranza, F. and M. Porreca (2012). Paleomagnetic secular variation at the Azores during the last 3 ka, *J. Geoph. Res.*, 117, B07101, DOI:10.1029/2012JB009285
50. C27n” transient greenhouse episode: evidences from marine pelagic Atlantic and peri-Tethyan sections. *Terra Nova*. doi: 10.1111/j.1365-3121.2012.01086.x
51. Festa V., Fiore A., Parisi M., Siniscalchi A (2012). Sinkhole Evolution in the Apulian Karst of Southern Italy: A Case Study, with Some Considerations on Sinkhole Hazards. *Journal Of Caves And Karst Studies*, vol. 74 (2), p. 137-147, ISSN: 1090-6924, doi: 10.4311/2011JCKS0211.
52. Galeotti, S., L. Lanci, F. Florindo, T. Naish, L. Sagnotti, S. Sandroni and F. Talarico (2012). Cyclochronology of the Eocene-Oligocene transition from a glaci-marine succession off the Victoria Land coast, Cape Roberts Project, Antarctica. *Palaeogeogr., Palaeoclimatol., Palaeocol.*, 335-336, 84-94, doi:10.1016/j.palaeo.2011.08.011
53. Jones, A. G., D. Kiyari, J. Fullea, J. Ledo, P. Queralt, A. Marcuello, Siniscalchi A, and G. Romano (2012). Comment on “Deep resistivity cross section of the intraplate Atlas Mountains (NW Africa): New evidence of anomalous mantle and related Quaternary volcanism”. *Tectonics*, vol. 31, p. 1-6, ISSN: 0278-7407, doi:10.1029/2011TC00305.
54. Lucifora, S., F. Cifelli, M. Mattei, L. Sagnotti, D. Cosentino and A.P. Roberts (2012), Inconsistent magnetic polarities in magnetite- and greigite-bearing sediments: understanding complex magnetizations in the late Messinian in the Adana basin (Southern Turkey), *Geochem. Geophys. Geosyst.* Vol. 13, No. 10, doi: 10.1029/2012GC004248.
55. Maffione M., S. Pucci, L. Sagnotti and F. Speranza (2012), Magnetic fabric of Pleistocene continental clays from the hanging-wall of an active low-angle normal fault (Altotiberina Fault, Italy), *Int. J. Earth Sci.* (Geol. Rundsch.), 101, 849-861, DOI 10.1007/s00531-011-0704-9
56. Mattei M., Cifelli F., Muttoni G., Zanchi A., Berra F., Mossavvari F., Safar Ali E. (2012), Neogene block-rotation in Central Iran: evidence from paleomagnetic data. *Geological Society American Bulletin*. doi:10.1130/B30479.1
57. Mercuri, A.M., Bandini Mazzanti M., Torri P., Vigliotti L., Bosi G., Florenzano A., Olmi L., Massamba N's I. 2012. A marine/terrestrial integration for mid-late Holocene vegetation history and the development of the cultural landscape in the Po valley as a result of human impact and climate change. *Veget Hist Archaeobot*; DOI 10.1007/s00334-012-0352-4
58. Nostro C., Amato A., Cultrera G., Margheriti L., Selvaggi G., Arcoraci L., Casarotti E., Di Stefano R., Cerrato S., May 11 Team (2012), Turning the rumor of the May 11, 2011, earthquake prediction in Rome, Italy, into an information day on earthquake hazard. *Annals of Geophysics*, 55, 3, doi: 10.4401/ag-5559, 2012.

59. Porreca, M. & Mattei, M. (2012) - AMS fabric and tectonic evolution of Quaternary intramontane extensional basins in the Picentini Mountains (Southern Apennines, Italy). *International Journal of Earth Sciences*. 101, 787–802, DOI: 10.1007/s00531-011-0670-2.
60. Roberts, A.P. Chang, L., Heslop, D., Florindo, F., Larrasoana, J.C. (2012) Searching for single domain magnetite in the ‘pseudo-single-domain’ sedimentary haystack: Implications of biogenic magnetite preservation for sediment magnetism and relative paleointensity determinations. *J. Geophys. Res.*, 117, B08104, doi:10.1029/2012JB009412.
61. Sagnotti, L. and A. Winkler (2012). On the magnetic characterization and quantification of the superparamagnetic fraction of traffic-related urban airborne PM in Rome, Italy, *Atmospheric Environment*, 59, 131-140.
62. Siniscalchi A., Tripaldi S., Neri M., Balasco M., Romano G., Ruch J., Schiavone D. (2012). Flank instability structure of Mt Etna inferred by a magnetotelluric survey. *Journal Of Geophysical Research, Solid Earth*, vol. 117, ISSN: 0148-0227, doi: 10.1029/2011JB008657.
63. Speranza, F., Di Chiara, A. and S. Rotolo (2012). Correlation of welded ignimbrites on Pantelleria (Strait of Sicily) using paleomagnetism, *Bull. Volcanol.*, 74, 341–357, DOI 10.1007/s00445-011-0521-9.
64. Speranza, F., L. Minelli, A. Pignatelli, and M. Chiappini (2012), The Ionian Sea: The oldest in situ ocean fragment of the world?, *J. Geophys. Res.*, 117, B12101, doi:10.1029/2012JB009475.
65. Vila, B., Galobart, À., Canudo, J. I., Le Loeuff, J., Dinarès-Turell, J., Riera, V., Oms, O., Tortosa, T., and Gaete, R. (2012). The diversity of sauropod dinosaurs and its first taxonomic succession from the latest Cretaceous strata of southwestern Europe: clues to demise and extinction. *Palaeogeogr., Palaeoclimatol., Palaeocol.*, 350-352, 19-38, doi:10.1016/j.palaeo.2012.06.008

## 2013

66. Alimohammadian H., Hamidi Z., Aslani A., Shahidi A., Cifelli F. & Mattei M. (2013). A tectonic origin of magnetic fabric in the Shemshak Group from Alborz Mts. (northern Iran). *Journal of Asian Earth Science*. doi.org/10.1016/j.jseaes.2013.05.014.
67. Angiolini L., Crippa G., Muttoni G. (2013). Guadalupian (Middle Permian) paleobiogeography of the Neotethys Ocean. *Gondwana Research*, 24, 173-184.
68. Cifelli F., Mattei M., Rashid H. & Ghalamghash J. (2013). Right-lateral transpressional tectonics along the boundary between Lut and Tabas blocks (Central Iran). *Geophys. J. Int.*, doi: 10.1093/gji/ggt070
69. Dall’Olio E., Felletti F., and Muttoni G. (2013). Magnetic-fabric analysis as a tool to constrain mechanism of deep-water mudstone deposition in the Marnoso-Arenacea Formation. *Journal of Sedimentary Research*, 83, 170-182.
70. De Marco, E., Tema, E., Lanos, Ph., Kondopoulou, D. (2013). An updated catalogue of Greek archaeomagnetic data and a directional secular variation curve for the last 4500 years. *Studia Geophysica Geodaetica*, 58, doi: 10.1007/s11200-013-0910-y.
71. De Santis A., Qamili E., Wu L.X., Toward a possible next geomagnetic transition ? *Nat. Haz. Earth Syst. Sc.*, 13, 3395-3403, 2013.
72. Dudkin F., Korepanov V., Hayakawa M., De Santis A., Possible model of electromagnetic signals before earthquakes, in *Thales* (ed. Arabelos D., Kaltsikis C., Spatalas S., Tziavos I.N.), 159-170, 2013.
73. Kent D.V., Muttoni G. (2013). Modulation of Late Cretaceous and Cenozoic climate by variable drawdown of atmospheric pCO<sub>2</sub> from weathering of basaltic provinces on continents drifting through the equatorial humid belt. *Climate of the Past*, 9, 525-546.
74. Lanci L. and B. Delmonte (2013) Magnetic properties of aerosol dust in peripheral and inner Antarctic ice cores as a proxy for dust provenance. *Global and Planetary Change*, 110, 414-419.
75. Lanci, L., E. Tohver, A. Wilson, S. Flint (2013) Upper Permian magnetic stratigraphy of the lower Beaufort Group, Karoo Basin, *Earth and Planetary Science Letters*, 375, 123–134.
76. Lucifora S., Cifelli F., Rojay F. B. & Mattei M. (2013) – Paleomagnetic rotations in the Late Miocene sequence from the Çankırı Basin (Central Anatolia, Turkey): the role of strike-slip tectonics. *Turkish Journal of Earth Sciences*, 22, doi:10.3906/yer-1207-2.

Qamili E., De Santis A., Isac A., Mandeia M., Duka B., Geomagnetic jerks as chaotic fluctuations of the Earth's magnetic field, *G-cubed*, vol.14, 839-850, 2013.

77. Muttoni, G., Dallanave, E., Channell, J.E.T., The drift history of Adria and Africa from 280 Ma to Present, Jurassic true polar wander, and zonal climate control on Tethyan sedimentary facies. *Paleogeography, Paleoclimatology, Paleogeography*, 386, 415-435.
78. Muttoni G., Scardia G., Kent, D.V. A critique of evidence for human occupation of Europe older than the Jaramillo Subchron (~1 Ma): Comment on "The oldest human fossil in Europe from Orce (Spain)" by Toro-Moyano et al. (2013). *Journal of Human Evolution*, 65, 746-749.
79. Singer B.S., Guillou H., Jicha B.R., Zanella E., Camps P., 2013, Refining the quaternary Geomagnetic Instability Time Scale (GITS): lava flow recordings of the Blake and post-Blake excursions. *Quaternary Geochronology*, <http://dx.doi.org/10.1016/j.quageo.2012.12.005>
80. Telesca L., Lovallo M., Romano G., Hsu H., Chiang C., Chen C. (2013), Informational analysis of apparent Earth's resistivity time series to assess the reliability of magnetotelluric measurements. *Journal of Asian Earth Sciences*, 77, 77–82
81. Tema, E., Fantino, F., Ferrara, E., Lo Giudice, A., Morales, J., Goguitchaichvili, A., Camps, P., Barelo, F., Gulmini, M. (2013). Combined archaeomagnetic and thermoluminescence study of a brick kiln excavated at Fontanetto Po (Vercelli, Northern Italy). *J. Arch. Science*, 40 (4), 2025-2035, <http://dx.doi.org/10.1016/j.jas.2012.12.011>.
82. Tema, E., Morales, J., Goguitchaichvili, A., Camps, P. (2013). New archaeointensity data from Italy and geomagnetic field intensity variation in the Italian Peninsula. *Geophysical Journal International*, 193 (2), 603-614.
83. Tema, E., Kondopoulou, D., Pavlides, S., (2013). Palaeotemperature estimation of the pyroclastic deposit covering the pre-Minoan palaeosol at Megalochory Quarry, Santorini (Greece): Evidence from magnetic measurements. *Studia Geophysica Geodaetica*, 57, doi: 10.1007/s11200-013-0601-8.
84. Tozzi R., De Santis A., Gaya-Piqué L.R., Antarctic geomagnetic reference model updated to 2010 and provisionally to 2012, *Tectonophysics*, 585,13-25. 2013.

## 2014

85. Agrò A., Zanella E., Le Pennec J.L., Temel A., (2014). Magnetic fabric of ignimbrites: a case study from the Central Anatolian Volcanic Province. Geological Society Special Publication, 396: "The Use of Palaeomagnetism and Rock Magnetism to Understand Volcanic Processes"
86. Aubele, K., Bachtadse, V., Muttoni, G., and Ronchi, A., 2014, Paleomagnetic data from Late Paleozoic dykes of Sardinia: Evidence for block rotations and implications for the intra-Pangea megashear system: Geochemistry, Geophysics, Geosystems. doi:10.1002/2014GC005325
87. Balasco M., Lapenna V., Romano G., Siniscalchi A., Stabile T.A., and Telesca L. (2014). Electric and Magnetic Field Changes Observed during a Seismic Swarm in Pollino Area (Southern Italy). *Bulletin Of The Seismological Society Of America*, vol. 104, ISSN: 0037-1106, doi: 10.1785/0120130183
88. Dallanave, E., C. Agnini, V. Bachtadse, G. Muttoni, J.S.Crampton, C. Percy Strong, B.R. Hines, C.J. Hollis, and B.S. Slotnick. Early to middle Eocene magneto-biochronology of the southwest Pacific Ocean and climate influence on sedimentation: Insights from the Mead Stream section, New Zealand. *Geological Society of America Bulletin*, doi: 10.1130/B31147.1
89. Di Chiara, A., F. Speranza, M. Porreca, A. Pimentel, F. D'Ajello Caracciolo, and J. Pacheco (2014), Constraining chronology and time-space evolution of Holocene volcanic activity on the Capelo Peninsula (Faial Island, Azores): The paleomagnetic contribution, *Geol. Soc. Am. Bull.*, 126, 1164–1180, doi:10.1130/B30933.1.
90. Di Chiara, A., L. Tauxe, and F. Speranza (2014), Paleointensity determination from São Miguel (Azores Archipelago) over the last 3 ka, *Phys. of the Earth and Planet. Int.*, 234,1-13.
91. Di Mauro, D., Alfonsi L., Sapia, V., Urbini, S. (2014). A neighborhood revealed by geophysical prospection: an example of urbanization at the Phoenician-Punic settlement of Mozia (western Sicily, Italy) . *Journal of Applied Geophysics*, vol. 104, p. 114-120, ISSN: 0926-9851
92. Dinarès-Turell J., Westerhold T., Pujalte V., Röhl, U. and Kroon D. (2014). Astronomical calibration of the



Danian Stage (Early Paleocene) revisited: settling chronologies across the Atlantic and Pacific Oceans. *Rend. Online Soc. Geol. It.*, Vol. 31, 64-65. doi: 10.3301/ROL.2014.47.

93. Dinarès-Turell J., Westerhold T., Pujalte V., Röhl, U. and Kroon D. (2014). Astronomical calibration of the Danian Stage (Early Paleocene) revisited: settling chronologies of sedimentary records across the Atlantic and Pacific Oceans. *Earth Planet. Sci. Lett.*, 405, 119-131. doi:10.1016/j.epsl.2014.08.027.
94. Dinarès-Turell, J., A. Payros, S. Monechi, X. Orue-Etxebarria, S. Ortiz, E. Apellaniz, and G. Bernaola (2014), In Search of the Bartonian (Middle Eocene) GSSP (II): Preliminary Results from the Oyambre Section (Northern Spain), in *STRATI 2013*, edited by R. Rocha, J. Pais, J. C. Kullberg and S. Finney, pp. 79-83, Springer International Publishing. DOI: 10.1007/978-3-319-04364-7\_16.
95. Dinarès-Turell, J., and Tinto, K., (2014). Data report: paleomagnetism and rock magnetism of sediments from offshore Canterbury Basin, IODP Expedition 317. In Fulthorpe, C.S., Hoyanagi, K., Blum, P., and the Expedition 317 Scientists, *Proc. IODP, 317: Tokyo (Integrated Ocean Drilling Program Management International, Inc.)*. DOI:10.2204/iodp.proc.317.206.2014.
96. Dinarès-Turell, J., T. Westerhold, V. Pujalte, U. Röhl, and D. Kroon (2014), Settling the Danian Astronomical Time Scale: A Prospective Global Unit Stratotype at Zumaia, Basque Basin, in *STRATI 2013*, edited by R. Rocha, J. Pais, J. C. Kullberg and S. Finney, pp. 191-195, Springer International Publishing. DOI: 10.1007/978-3-319-04364-7\_38
97. Dinarès-Turell, J., V. Pujalte, K. Stoykova, and J. Elorza (2014), Extending Back the Palaeogene Astronomical Time Scale: An Integrated Analysis of the Upper Maastrichtian Strata in the Basque Basin, in *STRATI 2013*, edited by R. Rocha, J. Pais, J. C. Kullberg and S. Finney, pp. 185-189, Springer International Publishing. DOI: 10.1007/978-3-319-04364-7\_37
98. Florindo, F., Bianco, F., De Michelis, P., Masina, S., Muscari, G., Caprara, F., Chiodetti, A.G., Del Pezzo, E., Gresta, S., *Annals of Geophysics: ad majora*. *Annals of Geophysics*, 57, doi:10.4401/ag-6532
99. Foresi L.M., Baldassini N., Sagnotti L., Lirer F., Di Stefano A., Caricchi C., Verducci M., Salvatorini G., Mazzei R. (2014), Integrated stratigraphy of the St. Thomas section (Malta Island): A reference section for the lower Burdigalian of the Mediterranean Region, *Marine Micropaleontology*, 111, 66-89.
100. Grant, K.M., Rohling, E.J., Bronk Ramsey, C., Cheng, H., Edwards, R.L., Florindo, F., Heslop, D., Marra, F., Roberts, A.P. Tamisiea, M.E., and Williams, F. (2014), Sea-level variability over five glacial cycles. *Nature Communications*, doi:10.1038/ncomms6076.
101. Hernandez-Moreno, C., F. Speranza, and A. Di Chiara (2014), Understanding kinematics of intraplate transcurrent deformation: Paleomagnetic evidence from the Liquiñe-Ofqui fault zone (Chile, 38-41°S), *Tectonics*, in press.
102. Macrì, P., Speranza, F., and Capraro, L., (2014). Magnetic fabric of Plio-Pleistocene sediments from the Crotone fore-arc basin: Insights on the recent tectonic evolution of the Calabrian Arc (Italy). *Journal of Geodynamics*. In press.
103. Marchetti M., Sapia V., Garello A., De Rita D., Venuti A. (2014) Geology and geophysics at the archeological park of Vulci (Central Italy), *Annals of Geophysics*, 57, 1, G0188; doi:10.4401/ag-6361.
104. Marra, F., Florindo, F., (2014), The subsurface geology of Rome, Italy. Relationships among sedimentary processes, sea-level changes and astronomical forcing. *Earth-Science Reviews*, <http://dx.doi.org/10.1016/j.earscirev.2014.05.001>
105. Mattei M., Riggs N., Conticelli S., Giordano G., Cifelli F., Soriano C., Jicha B., Marchionni S., Guarnieri L., Tommasini S., Jasim A., Franciosi L. & Porreca M. (in press). Geochronology Geochemistry and Geodynamics of the Cabo de Gata volcanic zone, South Eastern Spain. *Italian Journal of Geosciences*.
106. Muttoni, G., Tartarotti, P., Chiari, M., Marieni, C., Dallanave, E., Kirscher, U., Rodelli, D. (2014). Paleolatitudes of Late Triassic radiolarian cherts from Argolis, Greece: insights on the paleogeography of the western Tethys. *Palaeogeography, Palaeoclimatology, Palaeoecology*.
107. Muttoni G., Mazza M., Mosher D., Katz M.E., Kent D.V., Balini M., A Middle–Late Triassic (Ladinian–Rhaetian) carbon and oxygen isotope record from the Tethyan Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 399 (2014) 246–259.
108. Negri, A., Amorosi A., Antonioli F., Bertini A., Florindo F., Lurcock P.C., Marabini S., Mastronuzzi G., Regattieri E., Rossi V., Scarponi D., Taviani M., Zanchetta G., Vai G.B. (2014). A potential global boundary

stratotype section and point (GSSP) for the Tarentian Stage, Upper Pleistocene, from the Taranto area (Italy): Results and future perspectives, *Quaternary International*, <http://dx.doi.org/10.1016/j.quaint.2014.08.057>

109. Occhipinti A., De Santis A, Massimo Maffei, Magnetoreception: an unavoidable step for plant evolution?, *Trends in Plant Science*, Vol. 19, No.1, 1-4, 2014.
110. Pavon-Carasco F.J., M. L. Osete, J. M. Torta, A. De Santis, A geomagnetic field model of the Holocene based on archaeomagnetic and lava flow data, *Earth Planet. Science Lett.*, 388, 98-109, 2014.
111. Payros, A., J. Dinarès-Turell, X. Orue-Etxebarria, S. Monechi, S. Ortiz, E. Apellaniz, and G. Bernaola (2014), In Search of the Bartonian (Middle Eocene) GSSP (I): Potential in the Basque–Cantabrian and Aquitanian Basins (Western Pyrenees), in *STRATI 2013*, edited by R. Rocha, J.Pais, J. C. Kullberg and S. Finney, pp. 131-135, Springer International Publishing. DOI: 10.1007/978-3-319-04364-7\_27
112. Porreca M, F. Cifelli, C. Soriano, G. Giordano, C. Romano, S. Conticelli, Mattei M., 2014. Hyaloclastite fragmentation below the glass transition: example from El Barronal submarine volcanic complex (Spain). *Geology*, v. 42, 1, 87–90, doi:10.1130/G34744.1.
113. Romano G., Balasco M., Lapenna V., Siniscalchi A., Telesca L. and Tripaldi S. (2014). On the sensitivity of long-term magnetotelluric monitoring in Southern Italy and source-dependent robust single station transfer function variability. *Geophysical Journal International*, Issn: 0956-540X, doi: 10.1093/gji/ggu083
114. Sagnotti L., Scardia G., Giaccio B., Liddicoat J.C., Nomade S., Renne P.R. and C.J. Sprain (2014), Extremely rapid directional change during Matuyama-Brunhes geomagnetic polarity reversal, *Geophysical Journal International*, 199 (2): 1110-1124, doi: 10.1093/gji/ggu287
115. Savian, J., Jovane, L., Frontalini, F., Trindade, R., Coccioni, R., Bohaty, S., Wilson, P., Florindo, F., Roberts, A.P., Rita Catanzariti, R., Enhanced primary productivity and magnetotactic bacterial production in response to middle Eocene warming in the Neo-Tethys Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology*, DOI: 10.1016/j.palaeo.2014.08.009.
116. Schettino, A., 2014. *Quantitative Plate Tectonics*, 403 pp., Springer, Berlin, ISBN 978-3-319-09134
117. Soriano C., Giordano G., Riggs N., Porreca M., Bonamico A., Iosimi D., Cifelli F., Mattei M., De Benedetti A., Guarnieri L., Marchionni S. (2014). Geologic map, volcanic stratigraphy and structure of the Cabo de Gata volcanic zone, Betic-Rif orogen, SE Spain. *Italian Journal of Geosciences*.
118. Speranza, F., and L. Minelli (2014), Ultra-thick Triassic dolomites control the rupture behavior of the central Apennine seismicity: Evidence from magnetic modeling of the L’Aquila fault zone, *J. Geophys. Res.*, 119, doi:10.1002/2014JB011199.
119. Sulpizio R., Zanella E., Macias J.L., Saucedo R. (2014). Deposit temperature of pyroclastic density currents emplaced during the El Chichon 1982 and Colima 1913 eruptions. *Geological Society Special Publication* 396: “The Use of Palaeomagnetism and Rock Magnetism to Understand Volcanic Processes”
120. Stockhecke, M., O. Kwiecien, L. Vigliotti, F. Anselmetti, J. Beer, N. Cagatay, J.E.T. Channell, R. Kipfer, T. Litt, N. Pickarski, M. Sturm. (2014). Chronostratigraphy of the 600,000 year old long continental record of Lake Van (Turkey). *Quat. Sci. Rev.*, 104, pp. 8–17. doi:10.1016/j.quascirev.2014.04.008
121. Teitler, L., Kupp, G., Warnke, D.A., Florindo, F., Filippelli, G.M., Evidence from the Southern Ocean for a long warm interglacial across Marine Isotope Stage 31. *EPSL*, <http://dx.doi.org/10.1016/j.epsl.2014.10.037>.
122. Telesca L., Lovallo M., Romano G., Konstantinou K. I., Hsu H-L., Chen C. (2014). Using the informational Fisher–Shannon method to investigate the influence of long term deformation processes on geoelectrical signals: An example from the Taiwan orogeny. *Physica A*, vol. 414, p. 340-351, ISSN: 0378-4371, doi: 10.1016/j.physa.2014.07.060.
123. Tripanera D., Porreca M., Ruch J., Pimentel A., Acocella V., Pacheco J., Salvatore M. (2014). Relationships between tectonics and magmatism in transtensive/transform setting: an example from Faial Island (Azores, Portugal). *The Geological Society of America Bulletin*, v. 126, n. 1, 164–181, doi:10.1130/B30758.1
124. Vigliotti, L., J.E.T. Channell, M. Stockhecke. (2014). Paleomagnetism of Lake Van sediments: chronology and paleoenvironment since 350 ka. *Quat. Sci. Rev.*, 104, pp. 18–29. doi:10.1016/j.quascirev.2014.09.028

125. Vigliotti L. (2014). Magnetic properties of the Campanian Ignimbrite and the marine Y5 tephra layer. Geological Society, London, Special Publications 396: The Use of Palaeomagnetism and Rock Magnetism to Understand Volcanic Processes. doi: 10.1144/SP396.1
126. Villa G., Fioroni C., Persico D., Roberts A., Florindo F., (2014), Middle Eocene to Late Oligocene Antarctic Glaciation/Deglaciation and Southern Ocean productivity. *Paleoceanography*, doi:10.1002/2013PA002518

## **DIVISION II: “Aeronomic Phenomena”**

### **2011**

1. Alfonsi, Lu., L. Spogli, G. De Franceschi, V. Romano, M. Aquino, A. Dodson, and C. N. Mitchell (2011), Bipolar climatology of GPS ionospheric scintillation at solar minimum, *Radio Sci.*, 46, RS0D05, doi:10.1029/2010RS004571.
2. Alfonsi Lu., De Franceschi G., Romano V., Bourdillon A., Le Huy M., GPS scintillations and TEC gradients at equatorial latitudes on April 2006, *Adv. Space Res., Advances in Space Research* 47 (2011) 1750–1757
3. Alfonsi Lu., De Franceschi G., Romano V., Bourdillon A., Le Huy M., GPS scintillations and TEC gradients at equatorial latitudes on April 2006, *Adv. Space Res., Advances in Space Research* 47 (2011) 1750–1757
4. Bolaji, O. S.; Adeniyi, J. O.; Radicella, S. M.; Doherty, P. H. Variability of total electron content over an equatorial West African station during low solar activity. *RADIO SCIENCE* 47, DOI: 10.1029/2011RS004812
5. De Michelis, P., G. Consolini, M. Materassi, R. Tozzi, “An information theory approach to storm-substorm relationship”, *Journal of Geophysical Research*, vol. 116, A08225, doi:10.1029/2011JA016535, 2011.
6. Kashcheyev, A.; Nava, B.; Radicella, S. M. 2012. Estimation of higher-order ionospheric errors in GNSS positioning using a realistic 3-D electron density model. *RADIO SCIENCE* 47, DOI: 10.1029/2011RS004976
7. Materassi, M., L. Ciraolo, G. Consolini, N. Smith, “Predictive Space Weather: an information theory approach”, *Advances in Space Research* 47 (2011), pp. 877-885, doi:10.1016/j.asr.2010.10.026
8. Mikhailov A.V. and L. Perrone, On the mechanism of seasonal/solar cycle NmF2 variations: A quantitative estimate of the main parameters contribution using ISR observations, *J. Geophys. Res.*, 116, A03319, doi:10.1029/2010JA016122, 2011.
9. Mikhailov A.V. and L. Perrone, On the mechanism of seasonal/solar cycle NmF2 variations: A quantitative estimate of the main parameters contribution using ISR observations, *J. Geophys. Res.*, 116, A03319, doi:10.1029/2010JA016122, 2011.
10. Moreno B., S. Radicella, M. C. de Lacy, M. Herraiz, G. Rodriguez-Caderot; On the effects of the ionospheric disturbances on precise point positioning at equatorial latitudes, *GPS Solutions* 15:381–390 DOI 10.1007/s10291-010-0197-1, 2011
11. Moreno B., S. Radicella, M. C. de Lacy, M. Herraiz, G. Rodriguez-Caderot; On the effects of the ionospheric disturbances on precise point positioning at equatorial latitudes, *GPS Solutions* 15:381–390 DOI 10.1007/s10291-010-0197-1, 2011
12. Nava B.; Radicella, S.M.; Azpilicueta, F.. Data ingestion into NeQuick 2. *Radio Sci.* 46, RS0D17., doi:10.1029/2010RS004635, 2011

13. Oladipo O.A., J.O. Adeniyi, S.M. Radicella, I.A. Adimula, Variability of the ionospheric electron density at fixed heights and validation of IRI-2007 profile's prediction at Ilorin, *Advances in Space Research* ,47, 496–505 2011
14. Perrone L., A. Mikhailov and L. Korsunova, FORMOSAT-3/COSMIC E-region observations and daytime foE at middle latitudes, *J. Geophys. Res.*, 116, doi:10.1029/2010JA016411, 2011
15. Pezzopane M., M. Pietrella, A. Pignatelli, B. Zolesi, and L. R. Cander. Assimilation of autoscaled data and regional and local ionospheric models as input sources for real-time 3-D International Reference Ionosphere modelling. *Radio Science* , VOL. 46, RS5009, doi:10.1029/2011RS004697, September 2011
16. Prikryl P., Spogli L., Jayachandran P. T., Kinrade J., Mitchell C. N., Ning B., Li G., Cilliers P. J., Terkildsen M., Danskin D. W., Spanswick E., Weatherwax A. T., Bristow W. A., Alfonsi Lu., De Franceschi G., Romano V., Ngwira C. M., Opperman B. D. L. (2011), Interhemispheric comparison of GPS phase scintillation at high latitudes during the magnetic-cloud-induced geomagnetic storm of 5-7 April 2010, *ANNALES GEOPHYSICAE*, vol. 29, p. 2287-2304, ISSN: 0992-7689, doi: 10.5194/angeo-29-2287-2011
17. Sánchez-Cano B., M. Herraiz, G. Rodríguez-Caderot, S. M. Radicella; Study of the ionosphere of Mars: application and limitations of the Chapman-layer model, *Highlights of Spanish Astrophysics VI, Proceedings of the IX Scientific Meeting of the Spanish Astronomical Society, Madrid* (2011)
18. Sreeja V V, Aquino M H O, Forte B, Elmas Z, Hancock C, De Franceschi G, Alfonsi L, Spogli L, Romano V, Bougard B, Galera Monico J F, Wernik A W, Sleewaegen J-M, Canto´ M, Ferreira Da Silva E (2011). Tackling ionospheric scintillation threat to GNSS in Latin America. *JOURNAL OF SPACE WEATHER AND SPACE CLIMATE*, vol. 1, ISSN: 2115-7251

## 2012

19. Azzarone A., C.Bianchi, M.Pezzopane, M.Pietrella, C.Scotto, and A. Settini. IONORT: A Windows software tool to calculate the HF ray tracing in the ionosphere., *Computers & Geosciences*. Vol 42, pp 57-63. doi: 10.1016/j.cageo.2012.02.008, Maggio 2012
20. Katamzi Z. T., N. D. Smith, C. N. Mitchell, P. Spalla, M. Materassi, “Statistical analysis of traveling ionospheric disturbances using TEC observations over more than one solar cycle”, *Journal of Atmospheric and Solar-Terrestrial Physics* vol. 74 January, 2012. p. 64-80.
21. Magdaleno, Sergio; Herraiz, Miguel; Radicella, Sandro M. 2012. Ionospheric Bubble Seeker: A Java Application to Detect and Characterize Ionospheric Plasma Depletion From GPS Data. *IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING* 50 1719-1727
22. Marradi, L.; Zin, A.; Zago, S.; Landenna, S.; Mangolini, E.; Catalano, V.; Notarpietro, R.; Radicella, S.M. 2012. Nava, B. *The ROSA Project for Radio-Occultation: Findings and Current Status, Proceedings of the ION GNSS 2012, September 2012, Nashville, Tennessee.*
23. Mikhailov A.V., L. Perrone, N. Smirnova, Two types of positive disturbances in the daytime mid-latitude F2-layer: Morphology and formation mechanisms, *Journal Atm. Space Terr.* in press, 2012
24. Mikhailov A.V., A. Belehaki, L. Perrone, B. Zolesi, and I. Tsagouri, Retrieval of thermospheric parameters from routine ionospheric observations: assessment of method's performance at mid-latitudes daytime hours, *Journal of Space Weather and Space Climate*, 2, A03, 2012.
25. Mikhailov A.V., A. Belehaki, L. Perrone, B. Zolesi, and I. Tsagouri, Retrieval of thermospheric parameters from routinely observed F2-layer Ne(h) profiles at the geomagnetic equator, *Journal of Space Weather and Space Climate*, in press, 2012.
26. Nava, B.; Radicella, S.M.; Lagrasta, S.; Greco, F. 2012. Using NeQuick to reconstruct the 3D Electron Density of the Ionosphere: benefits and capabilities in single frequency positioning applications,

Proceedings of the IEEE First AESS European Conference on Satellite Telecommunications, October 2012, Rome, Italy.

27. Pietrella M., A short-term ionospheric forecasting empirical regional model (IFERM) to predict the critical frequency of the F2 layer during moderate, disturbed, and very disturbed geomagnetic conditions over the European area, *Ann. Geophys.*, 30, 343-355, doi: 10.5194/angeo-30-343-2012, Febbraio 2012.
28. Pietrella M., M. Pezzopane, C. Scotto. Variability of foF2 over Rome and Gibilmanna during three solar cycles (1976-2000), *J. Geophys. Res.*, Vol 117, A05316, doi:10.1029/2011JA017462, Maggio 2012
29. Katamzi Z. T., N. D. Smith, C. N. Mitchell, P. Spalla, M. Materassi, "Statistical analysis of traveling ionospheric disturbances using TEC observations over more than one solar cycle", *Journal of Atmospheric and Solar-Terrestrial Physics* vol. 74 January, 2012. p. 64-80.
30. Magdaleno, Sergio; Herraiz, Miguel; Radicella, Sandro M. 2012. Ionospheric Bubble Seeker: A Java Application to Detect and Characterize Ionospheric Plasma Depletion From GPS Data. *IEEE TRANSACTIONS ON GEOSCIENCE AND REMOTE SENSING* 50 1719-1727
31. Marradi, L.; Zin, A.; Zago, S.; Landenna, S.; Mangolini, E.; Catalano, V.; Notarpietro, R.; Radicella, S.M. 2012. Nava, B. *The ROSA Project for Radio-Occultation: Findings and Current Status, Proceedings of the ION GNSS 2012*, September 2012, Nashville, Tennessee.
32. Mikhailov A.V., L. Perrone, N. Smirnova, Two types of positive disturbances in the daytime mid-latitude F2-layer: Morphology and formation mechanisms, *Journal Atm. Space Terr.* in press, 2012
33. Mikhailov A.V., A. Belehaki, L. Perrone, B. Zolesi, and I. Tsagouri, Retrieval of thermospheric parameters from routine ionospheric observations: assessment of method's performance at mid-latitudes daytime hours, *Journal of Space Weather and Space Climate*, 2, A03, 2012.
34. Nava, B.; Radicella, S.M.; Lagrasta, S.; Greco, F. 2012. Using NeQuick to reconstruct the 3D Electron Density of the Ionosphere: benefits and capabilities in single frequency positioning applications, Proceedings of the IEEE First AESS European Conference on Satellite Telecommunications, October 2012, Rome, Italy.
35. Nigussie, M.; Radicella, S. M.; Damtie, B.; Nava, B.; Yizengaw, E.; Ciruolo, L. 2012. TEC ingestion into NeQuick 2 to model the East African equatorial ionosphere. *RADIO SCIENCE* 47, DOI: 10.1029/2012RS004981.
36. Pietrella M., A short-term ionospheric forecasting empirical regional model (IFERM) to predict the critical frequency of the F2 layer during moderate, disturbed, and very disturbed geomagnetic conditions over the European area, *Ann. Geophys.*, 30, 343-355, doi: 10.5194/angeo-30-343-2012, Febbraio 2012.
37. Pietrella M., M. Pezzopane, C. Scotto. Variability of foF2 over Rome and Gibilmanna during three solar cycles (1976-2000), *J. Geophys. Res.*, Vol 117, A05316, doi:10.1029/2011JA017462, Maggio 2012
38. Adewale, O.; Migoya Orué, Y. O. E.; Oyeyemi, O. Assessment of the NeQuick model at Lagos, Nigeria using GPS TEC, 2012, UNILAG GOLDEN JUBILEE RESEARCH CONFERENCE & FAIR, Proceedings Vol. III.
39. Nigussie, M.; Radicella, S. M.; Damtie, B.; Nava, B.; Yizengaw, E.; Ciruolo, L. 2012. TEC ingestion into NeQuick 2 to model the East African equatorial ionosphere. *RADIO SCIENCE* 47, DOI: 10.1029/2012RS004981.
40. Adewale, O.; Migoya Orué, Y. O. E.; Oyeyemi, O. Assessment of the NeQuick model at Lagos, Nigeria using GPS TEC, 2012, UNILAG GOLDEN JUBILEE RESEARCH CONFERENCE & FAIR, Proceedings Vol. III.

41. Alfonsi, Lu., L. Spogli, M. Pezzopane, V. Romano, E. Zuccheretti, G. De Franceschi, M. A. Cabrera, and R. G. Ezquer (2013), Comparative analysis of spread-F signature and GPS scintillation occurrences at Tucuman, Argentina, *J. Geophys. Res. Space Physics*, 118, doi:10.1002/jgra.50378.
42. Baumgardner J, J.Wroten, M. Mendillo, C.Martinis, C. Barbieri, G. Umbriaco, C. Mitchell, J. Kinrade, M. Materassi, L.Ciraolo and M. Hairston, "Imaging space weather over Europe", *SPACE WEATHER*, VOL. 11, 1–10, doi:10.1002/swe.20027, 2013.
43. Barta V., C. Scotto, M. Pietrella, V. Sgrigna, L. Conti, and G. Sàtori. *A statistical analysis on the relationship between thunderstorms and the sporadic E Layer over Rome*, *Astron.Nachr. /AN* 334, No. 9, 968 – 971 (2013) / DOI 10.1002/asna.201211972, November 2013.
44. Bianchi C, J.A. Baskaradas, M. Pezzopane, M.Pietrella, U.Sciacca, E. Zuccheretti. *Fading in the HF ionospheric channel and the role of irregularities*, *Advances in Space Research*, Vol 52, issue 3, pp 403-411, doi: 10.1016/j.asr.2013.03.035, August 2013.
45. Kutiev, I. Tsagouri, L. Perrone, D. Pancheva, P. Mukhtarov, A. Mikhailov ,J. Lastovicka, N. Jakowski, D. Buresova, E. Blanch, B. Andonov, D. Altadill, S. Magdaleno, M. Parisi, and J. Miquel Torta, Solar activity impact on the Earth's upper atmosphere, *Journal of Space Weather and Space Climate*, 3, A06, 2013.
46. Moen, J., Oksavik, K., Alfonsi, L., Daabakk, Y., Romano, V., & Spogli, L. (2013). Space weather challenges of the polar cap ionosphere. *Journal of Space Weather and Space Climate*, 3, A02.
47. Pietrella M.. Empirical regional models for the short-term forecast of M3000F2 during not quiet geomagnetic conditions over Europe, *Ann. Geophys.*, 31, 1653-1671, doi:10.5194/angeo-31-1653-2013, October 2013.
48. Romano V., Pau S., Pezzopane M., Spogli L., Zuccheretti E., Aquino M., Hancock C.M., eSWua: a tool to manage and access GNSS ionospheric data from mid-to-high latitudes, *Annals of Geophysics*, 56, 2, 2013, R0223; doi:10.4401/ag-6244
49. Romano V., Macelloni G., Spogli L., Brogioni M., Marinaro G., Mitchell C. N., Measuring GNSS ionospheric total electron content at Concordia, and application to L-band radiometers, *Annals of Geophysics*, 56, 2, 2013, R0219; doi:10.4401/ag-6241
50. Settini A., M. Pezzopane, M. Pietrella, C. Bianchi, C. Scotto, E. Zuccheretti, and J. Makris. *Testing the IONORT-ISP system: A comparison between synthesized and measured oblique ionograms*, *Radio Science*, VOL. 48(2), pp 1-13, doi:10.1002/rds.20018, April 2013.
51. Spogli L., Alfonsi L., Cilliers P.J., Correia E., De Franceschi G., Mitchell C.N., Romano V., Kinrade J., Cabrera M.A., GPS scintillations and total electron content climatology in the southern low, middle and high latitude regions, *Annals of Geophysics*, 56, 2, 2013, R0220; doi:10.4401/ag-6240
52. Spogli, L., Alfonsi, L., Romano, V., De Franceschi, G., Joao Francisco, G. M., Hirokazu Shimabukuro, M., ... & Aquino, M. (2013). Assessing the GNSS scintillation climate over Brazil under increasing solar activity. *Journal of Atmospheric and Solar-Terrestrial Physics*, 105, 199-206.
53. Tsagouri I. , A. Belehaki, N. Bergeot, C. Cid, V. Delouille, T. Egorova, N. Jakowski, I. Kutiev, A. Mikhailov, M. Nunez, M. Pietrella, A. Potapov, R. Qahwaji, Y. Tulunay, P. Velinov, and A. Viljanen. *Progress in space weather modeling in an operational environment*, *J. Space Weather Space Clim.* 3, A17 doi: 10.1051/swsc/2013037, April 2013.
54. Zolesi B. and Lj.R.Cander: *Ionospheric Prediction and Forecasting*, Springer and Verlag, Heidelberg, 2013.

2014

55. Baskaradas J. A., S. Bianchi, M. Pietrella, M. Pezzopane, U. Sciacca, E. Zuccheretti *Description of ionospheric disturbances observed by Vertical Ionospheric Sounding at 3 MHz*, Annals of Geophysics, 57, 1, 2014, A0187; doi:10.4401/ag-6345
56. Ezquer R. G., J. L. López, L. A. Scidá, M. A. Cabrera, B. Zolesi, C. Bianchi, M. Pezzopane, E. Zuccheretti, M. Mosert (2014), Behaviour of ionospheric magnitudes of F2 region over Tucumán during a deep solar minimum and comparison with the IRI 2012 model predictions, *Journal of Atmospheric and Solar Terrestrial Physics*, 107, 89-98, <http://dx.doi.org/10.1016/j.jastp.2013.11.010>.
57. Mikhailov A., A. Belehaki, L. Perrone, B. Zolesi, and I. Tsagouri, "Whether radio occultation middle latitude electron density profiles can be used to retrieve thermospheric parameters?", *Journal of Space Weather and Space Climate*, 4, 2014 <http://dx.doi.org/10.1051/swsc/20140092014>.
58. Mikhailov A. and L. Perrone, A method for foF2 short-term (1-24)h forecast using both historical and real-time foF2 observations over European stations: EUROMAP model, *Radio Science*, 2014, Accepted manuscript online: 23 MAR 2014 09:26PM EST | DOI: 10.1002/2014RS005373
59. Mikhailov A. and L. Perrone, Comment on "The winter anomaly in the middle-latitude F region during the solar minimum period observed by the Constellation Observing System for Meteorology, Ionosphere, and Climate" by Lee, W.K., Kil, H., Kwak, Y.-S., Wu, Q., Cho, S., and Park, J.U., *Journal of Geophysical Research*, 119, doi:10.1002/2014JA020185.2014.
60. Pezzopane M. (2014), Unusual nighttime impulsive enhancements of electron density characterizing the low-latitude ionosphere: Phenomenology and possible mechanisms of triggering, *Il Nuovo Cimento C* 37(4), 145-154, doi:10.1393/ncc/i2014-11799-9.
61. Pietrella M., M. Pezzopane, and C. Bianchi, *A comparative sporadic-E layer study between two mid-latitude ionospheric stations*, *Advances in Space Research*. Vol. 54 issue 2, pp 150-160, doi: DOI: 10.1016/j.asr.2014.03.019 , July 2014.
62. Pignalberi A., M. Pezzopane, E. Zuccheretti (2014), Sporadic E layer at mid-latitudes: average properties and influence of atmospheric tides, *Annales Geophysicae* 32, 1427-1440, doi:10.5194/angeo-32-1427-2014.
63. Settimi A., M. Pietrella, M. Pezzopane, B. Zolesi, C. Bianchi, and C. Scotto. The COMPLEIK subroutine of the IONORT-ISP system for calculating the non-deviative absorption: A comparison with the ICEPAC formula, *Advances in Space Research*. Vol 53, 2, Pages 201-218, January 2014. *Advances in Space Research*. Vol. 54 issue 2, pp 133-149, doi: DOI: 10.1016/j.asr.2014.03.018 , July, 2014

### Division III:

### "Magnetospheric Phenomena"

2011

1. Coco, I., G. Consolini, E. Amata, M.F. Marcucci, D. Ambrosino, Dynamical changes of the polar cap potential structure: an information theory approach, *Nonlinear Proc. Geophys.*, 18, 607-707, 2011
2. Consolini, G., P. De Michelis, Rank-ordering multifractal analysis of the auroral electrojet index, *Nonlinear Proc. Geophys.*, 18, 277-285, 2011.
3. De Michelis, P., G. Consolini, M. Materassi, R. Tozzi, An information theory approach to the storm-substorm relationship, *J. Geophys Res.*, 116, A08225, doi: 10.1029/2011JA016535, 2011

4. Di Lorenzo, C, P. Palangio, G. Santarato, A. Meloni, U. Villante, L. Santarelli, Non-inductive components of electromagnetic signals associated with L'Aquila earthquake sequences estimated by means of inter-station impulse response functions, *Nat. Hazards Earth Syst. Sci.*, doi:10.5194/nhess-11-1047-2011
5. Francia, P., M. Regi, M. De Lauretis, U. Villante, V.A. Pilipenko, A case study of upstream wave transmission to the ground at polar and low latitudes, *J. Geophys. Res.*, doi:10.1029/2011JA016751, in press, 2011.
6. Lepidi, S., L. Cafarella, M. Pietrolungo, L. Santarelli, Azimuthal propagation of Pc5 geomagnetic field pulsations in the southern polar cap, *Adv. Space Res.*, 47, 966-977, 2011
7. Lepidi, S., L. Cafarella, M. Pietrolungo, D. Di Mauro, Daily variation characteristics at polar geomagnetic observatories, *Adv. Space Res.*, 48, 521-528, 2011
8. Lepidi, S., A. Meloni, P. Palangio, K. Yumoto, The contribution of L'Aquila (Italy) Geomagnetic Observatory to MAGDAS project, *Sun and Geosphere*, 6, 49-53, 2011
9. Materassi, M., L. Ciraolo, G. Consolini, N. Smith, Predictive Space Weather: An information theory approach, *Adv. Space Res.*, 47, 877-885, 2011
10. Nenovski, P, M. Chamati, U. Villante, M. De Lauretis and P. Francia, Scaling characteristics of SEGMA magnetic field data around the Mw6.3 Aquila EQ, sub. "Statistical Mechanics in Earth Physics and Natural Hazards", *Acta Geophysica*, special issue, 2011.
11. Piersanti, M., U. Villante, C. Waters, I. Coco, The June 8, 2000 ULF wave activity: a case study, *J. Geophys. Res.*, doi:10.1029/2011JA016857, in press, 2011.
12. Prattes, G., K. Schwingenschuh, H. U. Eichelberger, W. Magnes, M. Boudjada, M. Stachel, M. Vellante, U. Villante, V. Wesztergom, P. Nenovski, Ultra Low Frequency (ULF) European multi station magnetic field analysis before and during the 2009 earthquake at L'Aquila regarding regional geotechnical information, *Nat. Hazards Earth Syst. Sci.*, 11, doi:10.5194/nhess,-11-1959-2011
13. Trenchi, L., Marcucci, M. F., Reme, H., Carr, C. M., Cao, J. B., TC-1 observations of a flux rope: Generation by multiple X line reconnection, *J. Geophys. Res.* 116, A5, A05202, 5202, 2011
14. Villante, U., C. De Paulis, P. Francia, The transmission of upstream waves to the magnetosphere: an analysis at widely separated ground stations, *J. Geophys. Res.*, 116, A06219, doi:10.1029/2010JA016263, 2011.
15. Villante, U., M. Piersanti, Sudden impulses at geosynchronous orbit and at low latitudes, *J. Atms. and Solar Terr. Phys.*, doi:10.1016/j.jastp.2010.01.008, 2011
16. Villante, U. and M. Piersanti, Sudden Impulses in the magnetosphere and at ground, in *The Solar Wind*, InTech Open Access Library, in press, 2011.

## 2012

17. Chamati, M., P. Nenovski, M. Vellante, U. Villante, K. Schwingenschuh, M. Boudjada, V. Wesztergom, Application of DFA method to magnetic field data, in *Complex Research of Earthquake's forecasting possibilities, Seismicity and Climate Change Correlations*, Ilia State University, Georgia, ISSN 2233-3681, pp. 72-79, 2012
18. De Michelis, P., G. Consolini, R. Tozzi, On the multi-scale nature of large geomagnetic storms: an empirical mode decomposition analysis, *Nonlinear Proc. Geophys.*, 19, 667-673, 2012



19. Francia P., M. Regi, M. De Lauretis, U. Villante, V.A. Pilipenko, A case study of upstream wave transmission to the ground at polar and low latitudes, *J. Geophys. Res.*, 117, A01210, doi:10.1029/2011JA016751, 2012.
20. Nenovski P., M. Chamati, U. Villante, M. De Lauretis, P. Francia, Scaling characteristics of SEGMA magnetic field data around the Mw 6.3 Aquila earthquake, *Acta Geophysica*, doi: 10.2478/s11600-012-0081-1, 2012.
21. Nenovski, P., M. Chamati, U. Villante, M. DeLauretis, M. Vellante, P. Francia, V. Wertztergom, K. Schwingenschuh, M. Boudjada, G. Prattes, DFA analysis of SEGMA magnetic field data around the M6.3 Aquila EQ, in *Complex Research of Earthquake's forecasting possibilities, Seismicity and Climate Change Correlations*, Ilia State University, Georgia, ISSN 2233-3681, pp. 157-163, 2012.
22. Piersanti, M., U. Villante, C. Waters, I. Coco, The June 8, 2000 ULF wave activity: a case study, *J. Geophys. Res.*, doi:10.1029/2011JA016857, 117, A02204, 2012.
23. Villante, U., M. Piersanti, Sudden Impulses in the magnetosphere and at ground, in *Exploring the Solar Wind*, InTech Open Access Library, 399, 2012

## 2013

24. Francia P., M. Regi, M. De Lauretis, ULF fluctuations observed along the SEGMA array during very low solar wind density conditions, *Planet. Space Sci.*, 81, 74-81, 2013.
25. Francia P., De Lauretis M, Regi M. (2013), ULF fluctuations observed along the SEGMA array during very low solar wind density conditions, *PLANETARY AND SPACE SCIENCE*, vol. 81, p. 74-81, ISSN: 0032-0633
26. Lichtenberger, J., M. A. Clilverd, B. Heilig, M. Vellante, J. Manninen, C. J. Rodger, A. B. Collier, A. M. Jørgensen, J. Reda, R. H. Holzworth, R. Friedel, and M. Simon-Wedlund (2013), The plasmasphere during a space weather event: first results from the PLASMON project, *J. Space Weather Space Clim.*, 3, A23, doi: <http://dx.doi.org/10.1051/swsc/2013045>.
27. Nenovski, P, M. Chamati, U. Villante, M. De Lauretis and P. Francia, Scaling characteristics of SEGMA magnetic field data around the Mw6.3 Aquila EQ, "Statistical Mechanics in Earth Physics and Natural Hazards", *Acta Geophysica*, 61, 2, 331, doi:10.2478/s11600-012-0081-1. 2013.
28. Nenovski P., M. Chamati, U. Villante, M. De Lauretis and P. Francia, Scaling characteristics of SEGMA magnetic field data around the Mw 6.3 Aquila earthquake, *Acta Geophysica*, 61, 311-337, doi: 10.2478/s11600-012-0081-1, 2013.
29. Regi, M., P. Francia, M. De Lauretis, K.-H. Glassmeier, U. Villante, Coherent transmission of upstream waves to polar latitudes through magnetotail lobes, *J. Geophys. Res.*, 118, 1-9, .., doi:10.1002/2012JA018472, 2013.
30. Villante, U. and M. Piersanti, On the Propagation of Sudden Impulses through the magnetosphere: the implications of a simple geometrical model, *J. Atm. Solar-Terr. Phys.*, [dx.doi.org/10.1016/j.jastp.2013.08.015](http://dx.doi.org/10.1016/j.jastp.2013.08.015), 2013.
31. Villante, U., Le Relazioni Sole-Terra e lo Space Weather, *Rendiconti Accademia Nazionale dei Lincei*, 2013 (in pubbl.).
32. Villante U., A. Del Corpo and P. Francia, Geomagnetic and solar wind fluctuations at discrete frequencies: a case study, *J. Geophys. Res.*, 118, 218-231, DOI: 10.1029/2012JA017971, 2013.

## 2014

33. Nenovski P. I., M. Pezzopane, L. Ciruolo, M. Vellante, U. Villante, De Lauretis M. (2014). Local changes in the total electron content immediately before the 2009 Abruzzo earthquake. *Advances in Space Research*, ISSN: 0273-1177, doi: 10.1016/j.asr.2014.09.029
34. Regi M., De Lauretis M., Francia P. (2014). The occurrence of upstream waves in relation with the solar wind parameters: a statistical approach to estimate the size of the foreshock region. *Planetary and space science*, vol. 90, p. 100-105, ISSN: 0032-0633
35. Regi M., De Lauretis M., Francia P., Villante U. (2014). The propagation of ULF waves from the Earth's foreshock region to ground: the case study of 15 February 2009. *Earth planets and space*, vol. 66:43, ISSN: 1343-8832, doi: 10.1186/1880-5981-66-43
36. Regi M., M. De Lauretis and P. Francia, The occurrence of upstream waves in relation with the solar wind parameters: a statistical approach to estimate the size of the foreshock region, *Planet. Space Sci.*, 90, 100-105, 2014.
37. Regi M., M. De Lauretis, P. Francia, U. Villante, The propagation of ULF waves from the Earth's foreshock region to ground: the case study of 15 February 2009, *Earth, Planets and Space*, 66:43, doi:10.1186/1880-5981-66-43, 2014.
38. Santarelli L, Palangio P., De Lauretis M. (2014). Electromagnetic background noise at L'Aquila Geomagnetic Observatory. *Annals of Geophysics*, vol. 57, ISSN: 1593-5213, doi: 10.4401/ag-6299
39. Vellante M., M. Piersanti, and E. Pietropaolo (2014), Comparison of equatorial plasma mass densities deduced from field line resonances observed at ground for dipole and IGRF models, *J. Geophys. Res. Space Physics*, 119, doi:10.1002/2013JA019568.
40. Villante, U., *Le Relazioni Sole-Terra e lo Space Weather*, Atti dei Convegni Lincei, Nazionale dei Lincei, 285, 103, 2014.
41. Villante, U., A. Piancatelli, and P. Palangio, On the man-made contamination on ULF measurements: evidence for disturbances related to an electrified DC railway, *Ann. Geophys.*, 32, 1153-1161, doi:10.5194/angeo-32-1153-2014, 2014.
42. Villante, U and P. Tiberi, A Comprehensive Analysis of the Occurrence and Characteristics of ULF Waves at Low Latitude, *J. Geophys. Res.* 2015, in press.
43. Villante, U. and M. Piersanti, Comment on "Statistical analysis of geosynchronous magnetic field perturbations near midnight during sudden commencements" by J.-S. Park et al., *sub. J. Geophys. Res.*, 2014.

#### **DIVISION IV:**

#### **"Solar Wind and Interplanetary Magnetic Field"**

#### **2011**

1. Alfonsi L., Spogli L, Giorgia De Franceschi, Vincenzo Romano, Marcio Aquino, Alan Dodson, Cathryn N. Mitchell (2011). Bipolar climatology of GPS ionospheric scintillation at solar minimum. In: *Proceeding of the IES2011: International Ionospheric Effects Symposium*.
2. D'Amicis R., R. Bruno, B. Bavassano, Response of the geomagnetic activity to solar wind turbulence during solar cycle 23, *Journal of Atmospheric and Solar-Terrestrial Physics*, 73, 653-657, 2011
3. Forte B., M. Materassi, Alfonsi Lu., V. Romano, G. De Franceschi, P. Spalla, Optimum parameter for estimating phase fluctuations on transionospheric signals at high latitudes, *J. Adv. Space Res.*, 47, 2188-2193, ISSN 0273-1177, DOI: 10.1016/j.asr.2010.04.033, 2011.

4. Gardini, M. Laurenza, M. Storini, SEP events and Multi-spacecraft Observations: Constraints on Theory, *Advances in Space Research*, 47, 2127-2139, 2011.
5. Marino R. et al., The magnetohydrodynamic turbulent cascade in the ecliptic solar wind: study of Ulysses data, *Plant. Space Sci.*, 59, 592, 2011
6. Perri, S., Carbone, V., Yordanova, E., Bruno, R., Balogh, A., Scaling law of the reduced magnetic helicity in fast streams. *Planetary and Space Science* 59, 575-579, 2011
7. Prikryl P, Spogli L, Jayachandran PT, Kinrade J, Mitchell CN, Ning B, Li G, Cilliers PJ, Terkildsen M, Danskin DW, Spanswick E, Donovan E, Weatherwax AT, Bristow WA, Alfonsi L, De Franceschi G, Romano V, Ngwira CM, Opperman BDL (2011). Interhemispheric comparison of GPS phase scintillation at high latitudes during the magnetic-cloud-induced geomagnetic storm of 5-7 April 2010. *ANNALES GEOPHYSICAE*, vol. 29, p. 2287-2304, ISSN: 0992-7689, doi: 10.5194/angeo-29-2287-2011.
8. Romano V. Bruno Bougard, Marcio Aquino, Joao F. Galera Monico, Tom Willems, Marc Solé, Investigation of low latitude scintillations in Brazil within the Cigala project, *Proceedings of the 3rd International Colloquium on Scientific and Fundamental Aspects of the Galileo Programme*, Copenhagen 2011, ESA Publications Division, European Space Agency, Noordwijk, The Netherlands, 2011.
9. Stangalini, M.; Del Moro, D.; Berrilli, F.; Jefferies, S. M., MHD wave transmission in the Sun's atmosphere, 2011 *A&A*...534A..65S
10. Viticchié, B.; Sánchez Almeida, J.; Del Moro, D.; Berrilli, F., Interpretation of HINODE SOT/SP asymmetric Stokes profiles observed in the quiet Sun network and internetwork, 2011 *A&A*...526A..60V

## 2012

11. Telloni, D., Bruno, R., D'Amicis, R., Pietropaolo, E., Carbone, V., Wavelet Analysis as a Tool to Localize Magnetic and Cross-helicity Events in the Solar Wind, accepted by *ApJ*\_751\_19 2012
12. Marino, R., Sorriso, L., Carbone, V., D'Amicis, R., Bruno, R., Veltri, P., On the nature of magnetohydrodynamic turbulence in high latitude solar wind, *ApJ*, 750\_41, 2012
13. D'Amicis, R., Consolini, G., Bavassano, B., Bruno, R., Conditioned Analysis Of High-Latitude Solar Wind Intermittency, *ApJ*, 755, 63, <http://dx.doi.org/10.1088/0004-637X/755/1/63>, 2012
14. Perri, S., Carbone, V., Vecchio, A., Bruno, R., Korth, H., Zurbuchen, T. H., and Sorriso, L., Phase-synchronization, Energy cascade and intermittency in the solar wind turbulence, *PRL*, 109, 245004, 2012
15. Antonucci, E. et al. including R. Bruno, Multi Element Telescope for Imaging and Spectroscopy (METIS) coronagraph for the Solar Orbiter mission, *Space Telescopes and Instrumentation 2012: Ultraviolet to Gamma Ray*, edited by Tadayuki Takahashi, Stephen S. Murray, Jan-Willem A. den Herder, *Proc. of SPIE* Vol. 8443, 844309, 2012
16. Owen et al. including R. Bruno, IMPALAS: Investigation of MagnetoPause Activity using Longitudinally-Aligned Satellites-a mission concept proposed for the ESA M3 2020/2022 launch. *Experimental Astronomy* 33, 365-401, 2012
17. M. Berthomier, et al. including R. Bruno, Alfvén: magnetosphere-ionosphere connection explorers, *Experimental Astronomy* 33, 445-489, 2012
18. Andretta V. et al., Coronal Diagnostics from Narrowband Images Around 30.4 nm, by., 2012, *Solar Physics*, 279, 53
19. Antonucci et al., UVCS Observations of Temperature and Velocity Profiles in Coronal Holes, 2012, *Space Science Review*, 172, 5

20. Macek, W.~M., Wawrzaszek, A., Carbone, V.\ 2012.\ Observation of the multifractal spectrum in the heliosphere and the heliosheath by Voyager 1 and 2.\ *Journal of Geophysical Research (Space Physics)* 117, 12101.
21. Carbone, V.\ 2012.\ Scalings, Cascade and Intermittency in Solar Wind Turbulence.\ *Space Science Reviews* 172, 343-360.
22. Lepreti, F., Carbone, V., Abramenko, V. I., Yurchyshyn, V., Goode, P. R., Capparelli, V., Vecchio, A.\ 2012.\ Turbulent Pair Dispersion of Photospheric Bright Points.\ *The Astrophysical Journal* 759, L17.
23. Donato, S., Servidio, S., Dmitruk, P., Carbone, V., Shay, M. A., Cassak, P. A., Matthaeus, W. H.\ 2012.\ Reconnection events in two-dimensional Hall magnetohydrodynamic turbulence.\ *Physics of Plasmas* 19, 092307.
24. Perri, S., Carbone, V.\ 2012.\ Comment on “Effect of Current Sheets on the Solar Wind Magnetic Field Power Spectrum from the Ulysses Observation: From Kraichnan to Kolmogorov Scaling”.\ *Physical Review Letters* 108, 199501.
25. Matthaeus, W. H., Servidio, S., Dmitruk, P., Carbone, V., Oughton, S., Wan, M., Osman, K. T.\ 2012.\ Local Anisotropy, Higher Order Statistics, and Turbulence Spectra.\ *The Astrophysical Journal* 750, 103.
26. Laurenza, M., Vecchio, A., Storini, M., Carbone, V.\ 2012.\ Quasi-biennial Modulation of Galactic Cosmic Rays.\ *The Astrophysical Journal* 749, 167.
27. Vecchio, A., Laurenza, M., Meduri, D., Carbone, V., Storini, M.\ 2012.\ The Dynamics of the Solar Magnetic Field: Polarity Reversals, Butterfly Diagram, and Quasi-biennial Oscillations.\ *The Astrophysical Journal* 749, 27.
28. A. Vecchio, M. Laurenza, M. Storini, V. Carbone, New Insights on Cosmic Ray Modulation through a Joint Use of Non stationary Data-Processing Methods, *Advances in Astronomy*, Volume 2012, Article ID 834247, doi:10.1155/2012/834247, 2012.
29. Romano, P.; Berrilli, F.; Criscuoli, S.; Del Moro, D.; Ermolli, I.; Giorgi, F.; Viticchié, B.; Zuccarello, F., A Comparative Analysis of Photospheric Bright Points in an Active Region and in the Quiet Sun, 2012 *SoPh*..280.407R
30. Giovannelli, L.; Berrilli, F.; Cocciolo, M.; Del Moro, D.; Egidi, A.; Piazzesi, R.; Stangalini, M., The birth of Tor Vergata Fabry-Pérot interferometer, 2012 *JPhCS*.383a2014G
31. Fabiani, S.; Bellazzini, R.; Berrilli, F.; Campana, R.; Costa, E.; Del Monte, E.; Muleri, F.; Rubini, A.; Soffitta, P., X-ray polarimetry towards high energy and solar science, 2012 *JPhCS*.383a2013F
32. Criscuoli, S.; Del Moro, D; Giannattasio, F.; Viticchié, B.; Giorgi, F.; Ermolli, I.; Zuccarello, F.; Berrilli, F., High cadence spectropolarimetry of moving magnetic features observed around a pore, 2012 *A&A*...546A..26C
33. Penza, V.; Berrilli, F., Velocity and Temperature Response Functions of 77 Near-Infrared (800 - 1400 nm) Photospheric Lines - I, 2012 *SoPh*..277..227P
34. Peter, H.; Abbo, L.; Andretta, V.; Auchère, F.; Bemporad, A.; Berrilli, F.; Bommier, V.; Braukhane, A.; Casini, R.; Curdt, W.; and 28 coauthors
35. Solar magnetism eXplorer (SolmeX). Exploring the magnetic field in the upper atmosphere of our closest star, 2012 *ExA*....33..271P
36. Stangalini, M.; Giannattasio, F.; Del Moro, D.; Berrilli, F., Three-minute wave enhancement in the solar photosphere, 2012 *A&A*...539L...4S

37. Sobotka, M.; Del Moro, D.; Jurčák, J.; Berrilli, F., Magnetic and velocity fields of a solar pore, 2012 A&A...537A..85S
38. A. Damiani, S. De Simone, C. Rafanelli, R. Cordero; M. Laurenza, Three years of ground-based total ozone measurements in Arctic: comparison with OMI and GOME satellites, Remote Sensing of Environment, 127, 162–180, 2012.
39. A. Vecchio, L. D'Alessi, V. Carbone, M. Laurenza, M. Storini, The Empirical Mode Decomposition to study the quasi-biennial modulation of solar magnetic activity and solar neutrino flux, Advances in Adaptive Data Analysis, Vol. 4, Nos. 1 & 2 (2012) 1250014 (17 pages), World Scientific Publishing Company, DOI: 10.1142/S1793536912500148.
40. C. Grimaldi, C. Boatella, M. Chmeissani, M. Fabi, N. Finetti, M. Laurenza, A. Lobo, I. Mateos, M. Storini, On the role of radiation monitors on board LISA Pathfinder and future space interferometers, Classical and Quantum Gravity, 29, 105001 (13pp), doi:10.1088/0264-9381/29/10/105001, 2012.
41. M. Laurenza, G. Consolini, M. Storini, A. Damiani, An information theory approach to the temporal evolution of SEP energy spectrum, Astrophysics and Space Sciences Transactions, 8, 19–24, 2012.

## 2013

42. Telloni, D., Perri, S., Bruno, R., Carbone, V., Amicis, R.~D. \ 2013. \ An Analysis of Magnetohydrodynamic Invariants of Magnetic Fluctuations within Interplanetary Flux Ropes. \ The Astrophysical Journal 776, 3.
43. Trenchi, L., Bruno, R., D'Amicis, R., Marcucci, M.~F., Telloni, D. \ 2013. \ Observations of IMF coherent structures and their relationship to SEP dropout events. \ Annales Geophysicae 31, 1333-1341.
44. Trenchi, L., Bruno, R., Telloni, D., D'Amicis, R., Marcucci, M.~F., Zurbuchen, T.~H., Weberg, M. \ 2013. \ Solar Energetic Particle Modulations Associated with Coherent Magnetic Structures. \ The Astrophysical Journal 770, 11.
45. Bruno, R., Carbone, V. \ 2013. \ The Solar Wind as a Turbulence Laboratory. \ Living Reviews in Solar Physics 10, 2.
46. Zank, G.~P., Bruno, R. and 12 colleagues 2013. \ SOLAR WIND 13: Proceedings of the Thirteenth International Solar Wind Conference. \ American Institute of Physics Conference Series 1539
47. Pucci, S., Poletto, G., Sterling, A.~C., Romoli, M. \ 2013. \ Physical Parameters of Standard and Blowout Jets. \ The Astrophysical Journal 776, 16.
48. Poletto, G., Sterling, A.~C., Pucci, S., Romoli, M. \ 2014. \ The contribution of X-ray polar blowout jets to the solar wind mass and energy. \ IAU Symposium 300, 239-242.
49. Pucci, S., Poletto, G., Sterling, A.~C., Romoli, M. \ 2013. \ An upper limit to the solar wind mass loading by X-ray polar jets. \ American Institute of Physics Conference Series 1539, 62-65.

## 2014

50. Bruno, R., Trenchi, L., Telloni, D. \ 2014. \ Spectral Slope Variation at Proton Scales from Fast to Slow Solar Wind. \ The Astrophysical Journal 793, LL15.
51. Bruno, R., Trenchi, L. \ 2014. \ Radial Dependence of the Frequency Break between Fluid and Kinetic Scales in the Solar Wind Fluctuations. \ The Astrophysical Journal 787, LL24.

52. Bruno, R., Telloni, D., Primavera, L., Pietropaolo, E., D'Amicis, R., Sorriso-Valvo, L., Carbone, V., Malara, F., Veltri, P.\ 2014.\ Radial Evolution of the Intermittency of Density Fluctuations in the Fast Solar Wind.\ *The Astrophysical Journal* 786, 53.
53. Zaqarashvili, T., Voeroes, Z., Narita, Y., Bruno, R.\ 2014.\ Twisted Magnetic Flux Tubes in the Solar Wind.\ *The Astrophysical Journal* 783, LL19.
54. Pucci, S., Poletto, G., Sterling, A.-C., Romoli, M.\ 2014.\ Birth, Life, and Death of a Solar Coronal Plume.\ *The Astrophysical Journal* 793, 86.
55. Poletto, G., Sterling, A.-C., Pucci, S., Romoli, M.\ 2014.\ The contribution of X-ray polar blowout jets to the solar wind mass and energy.\ *IAU Symposium* 300, 239-242.

## **DIVISION V:**

### **“Geomagnetic Observatories, Surveys, and Analyses ”**

#### **2011**

1. Barraclough D.R., De Santis A. , Repeat stations activities, in *Geomagnetic Observations and Models*, M. Manda and M. Korte (Ed.s), IAGA Special Sopron Book Series, vol. 5, 45-55, Springer, 2011..
2. Cafarella, L., S. Lepidi, A. Meloni, P. Palangio, M. Pietrolungo, L. Santarelli, *Geomagnetic Observation Results 2007-2008*, Terra Nova Bay, Antarctica, National Antarctic Research Program, PNRA, 24pp., 2011.
3. Cafarella, L., S. Lepidi, A. Meloni, P. Palangio, M. Pietrolungo, L. Santarelli, J. A. Baskaradas, *Geomagnetic Observation Results 2008-2009*, Terra Nova Bay, Antarctica, National Antarctic Research Program, PNRA, 24pp., 2011.
4. Cafarella, L., S. Lepidi, A. Meloni, P. Palangio, M. Pietrolungo, L. Santarelli, J. A. Baskaradas, , *Geomagnetic Observation Results 2009-2010*, Terra Nova Bay National Antarctica, Antarctic Research Program, PNRA, 25pp., 2011.
5. Di Lorenzo C., P. Palangio, G. Santarato, A. Meloni, U. Villante, and L. Santarelli, (2011) ‘Non-inductive components of electromagnetic signals associated with L’Aquila earthquake sequences estimated by means of inter-station impulse response functions’, *Nat. Hazards Earth Syst. Sci.*, 11, 1047–1055.
6. Lepidi S., L. Cafarella , M. Pietrolungo, D. Di Mauro, Daily variation characteristics at polar geomagnetic observatories, *Adv. Space Res.*, 48, 521-528, 2011.
7. Lepidi S., L. Cafarella , M. Pietrolungo, L. Santarelli, Azimuthal propagation of Pc5 geomagnetic field pulsations in the southern polar cap, *Adv. Space Res.*, 47, 966-977, 2011.
8. Lepidi S., A. Meloni, P. Palangio, K. Yumoto, (2011). The contribution of L'Aquila (Italy) Geomagnetic Observatory to MAGDAS project, *Sun and Geosphere*, 5, 17-22.

#### **2012**

9. Cafarella, L., S. Lepidi, A. Meloni, P. Palangio, M. Pietrolungo, L. Santarelli, J. A. Baskaradas, *Geomagnetic Observation Results 2010-2011*, Terra Nova Bay, Antarctica, National Antarctic Research Program, PNRA, 26pp., 2012.
10. Cianchini G., A. De Santis, D. R. Barraclough, L. X. Wu, and K. Qin, Magnetic transfer function entropy and the 2009 Mw = 6.3 L'Aquila earthquake (Central Italy), *Nonlin. Processes Geophys.*, 19, 2012, pp. 401-409, doi:10.5194/npg-19-401-2012

11. El Hawary R, K. Yumoto, Y. Yamazaki, A. Mahrous, E. Ghamry, A. Meloni, K. Badi, G. Kianji, C.B.S.Uiso, N. Mwiinga, L. Joao, Thomas Affluo, P. R. Sutcliffe, G. Mengistu, P. Baki, S.Abe, A.Ikeda, A. Fujimoto and T. Tokunaga, (2012). Annual and semi-annual Sq variations at 96 MM MAGDAS I and II stations in Africa, *Earth Planets Space*, 64, 425–432.
12. Tozzi R., G. Dominici, P. De Michelis, A. Meloni: Influence of solar activity on magnetic network measurements, *Annals of Geophysics*, 55, 6, 2012; doi: 10.4401/ag-5458.
13. Dominici G., A. Meloni, A. Di Ponzio, M. Miconi: Italian Magnetic Network and magnetic reference fields at 2010.0, *Annals of Geophysics*, 55, 6, 2012; doi: 10.4401/ag-5411.
14. Duka B., De Santis A., Manda M., Isac A., and E. Qamili, Geomagnetic jerks characterization via spectral analysis, *Solid Earth*, 3, 131–148, 2012.
15. Gasperini L., Cocchi L., Stanghellini C., Stanghellini G., Del Bianco F., Serrazanetti M., Carmisciano C., Magnetic and seismic-reflection study of Lake Cheko, a possible impact crater for the 1908 Tunguska Event. *Geochem. Geophys. Geosyst AGU* (2012), Volume 13, Number 5, 12 May 2012, Q05008, doi:10.1029/2012GC004054, ISSN: 1525-2027

## 2013

16. Palangio P., Carmisciano C., Di Lorenzo C., Caratterizzazione del rumore magnetico di fondo nel nuovo osservatorio di Varese Ligure. *Quaderni di Geofisica*, Anno 2013, Numero 109, ISSN 1590-2595.
17. Tozzi R., De Santis A., Gaya-Piqué L.R., Antarctic geomagnetic reference model updated to 2010 and provisionally to 2012, *Tectonophysics*, 2013, 585, 13-25.